

=> FILE REG

FILE 'REGISTRY' ENTERED AT 15:06:49 ON 23 APR 2010
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2010 American Chemical Society (ACS)

=> DISPLAY HISTORY FULL L1-

FILE 'HCA' ENTERED AT 11:43:49 ON 23 APR 2010
L1 16889 SEA SMITH D?/AU
L2 55459 SEA NITRIC#/TI AND OXIDE#/TI
L3 53 SEA L1 AND L2
L4 260698 SEA STABIL?/TI
L5 1 SEA L3 AND L4
SEL RN

FILE 'REGISTRY' ENTERED AT 11:46:03 ON 23 APR 2010
L6 14 SEA (10102-43-9/BI OR 113-21-3/BI OR 126-44-3/BI OR
E C9 H19 N5 O4 . NA/MF
L7 1 SEA "C9 H19 N5 O4 . NA"/MF
E DIAZENIUMDIOLATE
L8 2 SEA DIAZENIUMDIOLATE/BI
L9 2 SEA DIAZENIUMDIOL/BI

FILE 'LREGISTRY' ENTERED AT 11:51:38 ON 23 APR 2010
L10 STR

FILE 'REGISTRY' ENTERED AT 12:04:24 ON 23 APR 2010
L11 50 SEA SSS SAM L10
L12 21596 SEA SSS FUL L10
SAV TEM L12 MAR753/A

FILE 'LREGISTRY' ENTERED AT 12:06:32 ON 23 APR 2010
L13 STR
L14 STR

FILE 'REGISTRY' ENTERED AT 12:26:14 ON 23 APR 2010
L15 50 SEA SUB=L12 SSS SAM L13 OR L14
L16 1329 SEA SUB=L12 SSS FUL L13 OR L14
SAV L16 MAR573A/A
E NITRIC OXIDE/CN
L17 1 SEA "NITRIC OXIDE"/CN
L18 545 SEA (N (L) O)/ELS (L) 2/ELC.SUB

FILE 'HCA' ENTERED AT 12:33:54 ON 23 APR 2010
L19 7156 SEA L17/P OR L18/P

FILE 'LCA' ENTERED AT 12:34:04 ON 23 APR 2010
L20 56 SEA (PRODUC? OR PROD# OR GENERAT? OR MANUF? OR MFR# OR
CREAT? OR FORM## OR FORMING# OR FORMAT? OR MAKE# OR MADE#

OR MAKING# OR FABRICAT? OR SYNTHESI? OR PREPAR? OR
 PREP#) (3A) ((NITRIC# OR NITROUS# OR NITROGEN# OR N) (A) (OXIDE
 # OR MONOXIDE# OR DIOXIDE# OR TRIOXIDE# OR TETRAOXIDE# OR
 TETROXIDE#))

L21 0 SEA (PRODUC? OR PROD# OR GENERAT? OR MANUF? OR MFR# OR
 CREAT? OR FORM## OR FORMING# OR FORMAT? OR MAKE# OR MADE#
 OR MAKING# OR FABRICAT? OR SYNTHESI? OR PREPAR? OR
 PREP#) (3A) ((NITRIC# OR NITROUS# OR NITROGEN# OR N) (A) (PENTO
 XIDE# OR PENTAOXIDE#))

L22 43 SEA (PRODUC? OR PROD# OR GENERAT? OR MANUF? OR MFR# OR
 CREAT? OR FORM## OR FORMING# OR FORMAT? OR MAKE# OR MADE#
 OR MAKING# OR FABRICAT? OR SYNTHESI? OR PREPAR? OR
 PREP#) (2A) (NOX OR NO2 OR NO4 OR NO5 OR N2O OR N2O2 OR N2O3
 OR N2O4 OR N2O5 OR N3O OR N3O2 OR N3O3 OR N3O4 OR N3O5)

L23 0 SEA (PRODUC? OR PROD# OR GENERAT? OR MANUF? OR MFR# OR
 CREAT? OR FORM## OR FORMING# OR FORMAT? OR MAKE# OR MADE#
 OR MAKING# OR FABRICAT? OR SYNTHESI? OR PREPAR? OR
 PREP#) (2A) (N4O OR N4O2 OR N4O3 OR N4O4 OR N4O5 OR N5O OR
 N5O2 OR N5O3 OR N5O4 OR N5O5)

FILE 'HCA' ENTERED AT 13:53:59 ON 23 APR 2010

L24 288268 SEA (ION OR IONS OR IONIC? OR CATION? OR ANION?) (2A) (EXCHAN
 G? OR INTERCHANG?)

L25 76352 SEA L20 OR L21 OR L22 OR L23

L26 675 SEA (L19 OR L25) AND L24

L27 1 SEA L26 AND L5

FILE 'REGISTRY' ENTERED AT 13:59:09 ON 23 APR 2010

L28 1 SEA 113-21-3

L29 1 SEA 126-44-3

L30 1 SEA 14265-44-2

L31 1 SEA 9000-11-7

L32 1 SEA 9003-53-6

L33 1 SEA 9004-34-6

L34 1 SEA 9012-76-4

L35 373 SEA DOWEX#

L36 1 SEA L35 AND L6

FILE 'HCA' ENTERED AT 14:00:14 ON 23 APR 2010

L37 2 SEA L36

L38 239076 SEA L35 OR DOWEX#

L39 1 SEA L26 AND L37

L40 25 SEA L26 AND L38

L41 64445 SEA ANION? (2A) (EXCHANG? OR INTERCHANG?)

L42 5 SEA L40 AND L41

L43 153 SEA L26 AND L41

L44 44 SEA L43 AND L19

FILE 'REGISTRY' ENTERED AT 14:06:04 ON 23 APR 2010

E ASCORBATE/CN

L45 1 SEA ASCORBATE/CN

E NITRITE/CN

L46

1 SEA NITRITE/CN


FILE 'HCA' ENTERED AT 14:06:47 ON 23 APR 2010

L47 39929 SEA L45 OR ASCORBATE#
L48 91467 SEA L46 OR NITRITE#
L49 114076 SEA L28 OR LACTATE#
L50 3 SEA L40 AND (L47 OR L48 OR L49)
L51 39 SEA L43 AND (L47 OR L48 OR L49)
L52 16 SEA L44 AND (L47 OR L48 OR L49)
L53 120113 SEA L17
L54 3425 SEA L17/P
L55 6 SEA L54 AND L41
L56 12 SEA L54 AND (L37 OR L38)
L57 17 SEA L54 AND L24
L58 6 SEA L57 AND (L47 OR L48 OR L49)
L59 1080 SEA L53 AND L24
L60 3 SEA L59 AND L47
L61 63 SEA L59 AND L48
L62 5 SEA L59 AND L49
L63 10250 SEA (CREAM? OR GEL OR GELS OR GELLED OR GELLING#) (3A)L24
L64 1 SEA L63 AND L54
L65 17 SEA L63 AND L53
L66 16 SEA L63 AND L25
L67 9 SEA L65 AND L66
L68 2 SEA L63 AND L19
L69 34 SEA L39 OR L42 OR L50 OR L55 OR L58 OR L60 OR L62 OR L67
OR L68
L70 32 SEA (L52 OR L56 OR L57) NOT L69
L71 17 SEA L40 NOT (L69 OR L70)
L72 26 SEA 1808-2003/PY,PRY,AY AND L69
L73 26 SEA 1808-2003/PY,PRY,AY AND L70
L74 15 SEA 1808-2003/PY,PRY,AY AND L71
L75 20 SEA L53 AND L72
L76 13 SEA L53 AND L73
L77 1 SEA L53 AND L74
L78 14 SEA L77 OR L76
L79 QUE NANO?
L80 230896 SEA L17 OR L18
L81 285 SEA ?DIAZENIUMDIOL?
L82 5549 SEA L16
L83 390 SEA (L80 OR L25) AND (L81 OR L82)
L84 32 SEA L83 AND L79
L85 78266 SEA L12
L86 36403 SEA (L80 OR L25) AND L85
L87 803 SEA L86 AND L79
L88 6016 SEA L31
L89 139276 SEA L32
L90 115124 SEA L33
L91 34303 SEA L34
L92 15 SEA L87 AND (L88 OR L89 OR L90 OR L91)
L93 814 SEA L28
L94 1749 SEA L29

L95 47972 SEA L30
 L96 3 SEA L87 AND (L93 OR L94 OR L95)
 L97 16 SEA (L19 OR L25) AND (L81 OR L82) AND L79
 L98 243 SEA (L19 OR L25) AND L85 AND L79
 L99 4 SEA L98 AND ((L88 OR L89 OR L90 OR L91))
 L100 2 SEA L98 AND ((L93 OR L94 OR L95))
 L101 31 SEA L92 OR L96 OR L97 OR L99 OR L100
 L102 16 SEA L84 NOT L101
 L103 19 SEA 1808-2003/PY,PRY,AY AND L101
 L104 6 SEA 1808-2003/PY,PRY,AY AND L102
 L105 33244 SEA NANO2
 L106 12 SEA L103 NOT L105
 L107 3 SEA L104 NOT L105

FILE 'REGISTRY' ENTERED AT 15:06:49 ON 23 APR 2010

=> D L16 QUE STAT
 L10 STR



NODE ATTRIBUTES:

HCOUNT IS E0 AT 4
 NSPEC IS RC AT 2
 CONNECT IS E2 RC AT 3
 CONNECT IS E1 RC AT 4
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 3

STEREO ATTRIBUTES: NONE

L12 21596 SEA FILE=REGISTRY SSS FUL L10
 L13 STR



NODE ATTRIBUTES:


DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 6

STEREO ATTRIBUTES: NONE
L14 STR



NODE ATTRIBUTES:
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
RING(S) ARE ISOLATED OR EMBEDDED
NUMBER OF NODES IS 5

STEREO ATTRIBUTES: NONE
L16 1329 SEA FILE=REGISTRY SUB=L12 SSS FUL L13 OR L14

100.0% PROCESSED 2439 ITERATIONS 1329 ANSWERS
SEARCH TIME: 00.00.01

=> FILE HCA
FILE 'HCA' ENTERED AT 15:07:15 ON 23 APR 2010
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2010 AMERICAN CHEMICAL SOCIETY (ACS)

CLAIM 18 AND RELATED

=> D L106 1-12 BIB ABS HITSTR HITIND

L106 ANSWER 1 OF 12 HCA COPYRIGHT 2010 ACS on STN
AN 143:80337 HCA Full-text
TI Systems for preparing fine particles and other substances
IN Iversen, Steen Brummerstedt; Felsvang, Karsten; Larsen, Tommy;
Luethje, Viggo
PA SCF Technologies A/S, Den.
SO PCT Int. Appl., 53 pp.

CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2005058472	A2	20050630	WO 2004-DK888	20041219
	AU 2004298723	A1	20050630	AU 2004-298723	20041219
	AU 2004298723	B2	20080710		
	CA 2550518	A1	20050630	CA 2004-2550518	20041219
	CA 2550518	C	20100209		
	EP 1699549	A2	20060913	EP 2004-803039	20041219
	CN 1909955	A	20070207	CN 2004-80040700	20041219
	JP 2007514529	T	20070607	JP 2006-544222	20041219
	IN 2006DN04056	A	20070713	IN 2006-DN4056	20060714
	KR 2006130612	A	20061219	KR 2006-714539	20060719
	US 20070265357	A1	20071115	US 2007-583024	20070322
PRAI	DK 2003-1899	A	20031219		
	WO 2004-DK888	W	20041219		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The controlled prepn. of fine particles such as ~~nano~~-cryst. films and powders with at least one solvent being in a supercrit. state is carried out by introducing substances dissolved and/or dispersed in a solvent into a vessel and allowing the substances to ppt. at least partly as primary particles on the surface or said material. Further treatment of formed particles such as encapsulation of formed primary particles and collection of formed substances in a batch wise, semi-continuous or continuous manner can be carried out.

IT 9003-53-6, Polystyrene 10024-97-2, Nitrous oxide, reactions

(systems for prepg. fine particles and other substances)


RN 9003-53-6 HCA

CN Benzene, ethenyl-, homopolymer (CA INDEX NAME)

CM 1


CRN 100-42-5

CMF C8 H8



RN 10024-97-2 HCA

CN Nitrogen oxide (N2O) (CA INDEX NAME)



IC ICM B01J002-00
 CC 48-8 (Unit Operations and Processes)
 ST ~~nanoparticle~~ manuf
 IT Aerogels
 Antibacterial agents
 Ceramics
 Drugs
 Encapsulation
 Ferromagnetic materials
 Magnetic materials
 Nanoparticles
 Paramagnetic materials
 Piezoelectric materials
 Sound and Ultrasound
 Surfactants
 Waters
 (systems for prepg. fine particles and other substances)
 IT 64-17-5, Ethanol, reactions 64-19-7, Acetic acid, reactions
 67-56-1, Methanol, reactions 67-63-0, Isopropanol, reactions
 67-64-1, Acetone, reactions 67-68-5, DMSO, reactions 71-23-8,
 Propanol, reactions 71-36-3, Butanol, reactions 71-41-0, Pentanol,
 reactions 74-82-8, Methane, reactions 74-84-0, Ethane, reactions
 74-85-1, Ethylene, reactions 74-98-6, Propane, reactions 75-72-9,
 Chlorotrifluoromethane 77-92-9, Citric acid, reactions 78-79-5D,
 Isoprene, polymers 78-83-1, Isobutanol, reactions 106-97-8,
 Butane, reactions 107-21-1, Ethylene glycol, reactions 109-66-0,
 Pentane, reactions 109-99-9, THF, reactions 110-54-3, Hexane,
 reactions 110-82-7, Cyclohexane, reactions 111-27-3, Hexanol,
 reactions 121-69-7, N,N-Dimethylaniline, reactions 124-38-9,
 Carbon dioxide, reactions 142-82-5, Heptane, reactions 593-53-3,
 Monofluoromethane 2551-62-4, Sulfur hexafluoride 7664-41-7,
 Ammonia, reactions 9002-86-2, Polyvinyl chloride 9002-88-4,
 Polyethylene 9003-05-8, Polyacrylamide 9003-07-0, Polypropylene
 9003-20-7, Polyvinyl acetate ~~9003-53-6~~, Polystyrene
~~10024-97-2~~, Nitrous oxide, reactions
 25038-59-9, reactions 25322-68-3, Polyethylene glycol
 (systems for ~~prepg.~~ fine particles and other substances)
 OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)
 RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

 L106 ANSWER 2 OF 12 HCA COPYRIGHT 2010 ACS on STN
 AN 143:32415 HCA Full-text
 TI Soft tissue implants and anti-scarring agents
 IN Hunter, William L.; Gravett, David M.; Toleikis, Philip M.; Maiti,
 Arpita
 PA Angiotech International A.-G., Switz.
 SO PCT Int. Appl., 2592 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 19

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2005051444	A2	20050609	WO 2004-US39465	20041122
	US 20050148512	A1	20050707	US 2004-986230	20041110
	US 20050181977	A1	20050818	US 2004-986231	20041110
	CN 101094613	A	20071226	CN 2004-80031664	20041110
	AU 2004293075	A1	20050609	AU 2004-293075	20041122
	CA 2536192	A1	20050609	CA 2004-2536192	20041122
	WO 2005051232	A2	20050609	WO 2004-US39346	20041122
	WO 2005051232	A3	20051208		
	WO 2006055008	A2	20060526	WO 2004-US39353	20041122
	WO 2006055008	A3	20090416		
	EP 1687041	A2	20060809	EP 2004-812062	20041122
	CN 1878514	A	20061213	CN 2004-80033341	20041122
	JP 2007514472	T	20070607	JP 2006-541689	20041122
	US 20050149158	A1	20050707	US 2004-409	20041129
	US 20050175662	A1	20050811	US 2004-451	20041129
	US 20050175661	A1	20050811	US 2004-999205	20041129
	US 20050186243	A1	20050825	US 2004-97	20041129
	US 20050186242	A1	20050825	US 2004-999204	20041129
	US 20050191331	A1	20050901	US 2004-1419	20041130
	US 20050175663	A1	20050811	US 2004-1791	20041202
	US 20050181008	A1	20050818	US 2004-1786	20041202
	US 20050181011	A1	20050818	US 2004-1792	20041202
	US 20050143817	A1	20050630	US 2004-6899	20041207
	US 20050177103	A1	20050811	US 2004-6314	20041207
	US 20050177225	A1	20050811	US 2004-6895	20041207
	US 20050181004	A1	20050818	US 2004-6289	20041207
	US 20060147492	A1	20060706	US 2006-343809	20060131
	ZA 2006002379	A	20091028	ZA 2006-2379	20060323
	CN 101420970	A	20090429	CN 2004-80033576	20060515
	IN 2006KN01694	A	20070511	IN 2006-KN1694	20060619
	IN 2006KN01695	A	20070511	IN 2006-KN1695	20060619
	IN 2006KN01698	A	20070511	IN 2006-KN1698	20060619
PRAI	US 2003-523908P	P	20031120		
	US 2003-524023P	P	20031120		
	US 2003-525226P	P	20031124		
	US 2003-526541P	P	20031203		
	US 2004-578471P	P	20040609		
	US 2004-586861P	P	20040709		
	US 2004-986230	A	20041110		
	US 2004-986231	A	20041110		
	US 2003-518785P	P	20031110		
	US 2004-582833P	P	20040624		
	US 2004-986450	A1	20041110		
	WO 2004-US37930	W	20041110		
	WO 2004-US39183	W	20041122		
	WO 2004-US39346	W	20041122		
	WO 2004-US39353	W	20041122		
	WO 2004-US39465	W	20041122		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The invention relates to soft tissue implants for use in cosmetic or reconstructive surgery and to compns. to make the implants resistant to growth by inflammatory scar tissue. Thus, a silicone gel contg. paclitaxel was used as a filling in breast implant.

IT 10102-43-9, Nitrogen oxide (NO), biological studies
(soft tissue implants and anti-scarring agents)

RN 10102-43-9 HCA


CN Nitrogen oxide (NO) (CA INDEX NAME)



IT 13010-20-3D, Nitrosourea, derivs.
(soft tissue implants and anti-scarring agents)

RN 13010-20-3 HCA

CN Urea, N-nitroso- (CA INDEX NAME)



IT 9012-76-4, Chitosan
(soft tissue implants and anti-scarring agents)

RN 9012-76-4 HCA

CN Chitosan (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

IC ICM A61L027-00
ICS A61L027-54; A61L031-00; A61L031-16

CC 63-7 (Pharmaceuticals)
Section cross-reference(s): 1, 62

IT Drug delivery systems
(nanospheres; soft tissue implants and anti-scarring agents)

IT 56-23-5, biological studies 10102-43-9, Nitrogen oxide (NO),
biological studies
(soft tissue implants and anti-scarring agents)

IT 50-07-7, Mitomycin C 50-44-2, 6-Mercaptopurine 51-21-8, 5-FU
53-79-2 55-21-0, Benzamide 55-86-7, Nitrogen mustard 57-22-7
59-05-2, Methotrexate 65-46-3D, Cytidine, analogs 69-33-0,
Tubercidin 98-92-0, Nicotinamide 107-41-5, Hexylene glycol
120-73-0D, Purine, analogs 127-07-1, Hydroxyurea 127-07-1D,
Hydroxyurea, derivs. 129-56-6, SP 600125 147-94-4, Cytarabine
289-95-2D, Pyrimidine, analogs 459-73-4, Ethyl glycine 501-36-0,
Resveratrol 518-28-5, Podophyllotoxin 865-21-4, Vinblastine
1404-15-5 3672-15-9, D-Mannose 6-phosphate 4291-63-8, Cladribine
7059-24-7, Chromomycin A3 7440-06-4D, Platinum, compds. 7689-03-4,
Camptothecin 7689-03-4D, Camptothecin, derivs. 7784-18-1, Aluminum

fluoride (AlF₃) 7789-20-0, Deuterium oxide 10540-29-1
~~13010-20-3D~~, Nitrosourea, derivs. 14110-64-6, Cytochalasin A
 15663-27-1, Cisplatin 18378-89-7 18457-55-1 19542-67-7, BAY
 11-7082 20830-81-3 22668-01-5 22862-76-6 23214-92-8
 24280-93-1, Mycophenolic acid 25316-40-9 25812-30-0 28128-19-0,
 2-Mercaptopurine 30562-34-6, Geldanamycin 31698-14-3 32222-06-3,
 1 α -25-Dihydroxyvitamin D₃ 33069-62-4 33419-42-0, Etoposide
 34031-32-8, Auranofin 34157-83-0 36877-68-6D, Nitroimidazole,
 derivs. 41859-67-0 52214-84-3 53123-88-9, Rapamycin
 53123-88-9D, Rapamycin, desmethyl derivs. 55837-20-2 56390-09-1
 58957-92-9 61318-90-9, Sulconazole 61825-94-3 64222-94-2,
 15-Deoxyprostaglandin J₂ 65271-80-9 70539-42-3 71486-22-1
 74913-06-7, Chromomycin 75330-75-5, Lovastatin 79902-63-9
 84625-61-6, Itraconazole 86160-53-4D, analogs 95058-81-4,
 Gemcitabine 98629-43-7, Gusperimus 104987-11-3, Tacrolimus
 114719-57-2 114977-28-5, Docetaxel 128794-94-5, Mycophenolate
 mofetil 137071-32-0, Pimecrolimus 149550-36-7, LY 290181
 152121-30-7, SB 202190 159351-69-6, Everolimus 160677-67-8,
 Tresperimus 164301-51-3, CNI 1493 173026-17-0, BXT 51072
 186692-46-6, CYC 202 189453-10-9 222036-17-1, GW 8510
 254750-02-2, IDN 6556 329773-35-5, Bay 58-2667 467214-20-6
 851536-75-9, Biolimus A₉

(soft tissue implants and anti-scarring agents)

IT 50-02-2, Dexamethasone 50-28-2, Estradiol, biological studies
 51-45-6, Histamine, biological studies 56-53-1, Diethyl stilbestrol
 57-50-1D, Sucrose, derivs. 62-55-5, Thioacetamide 64-17-5,
 Ethanol, biological studies 79-10-7D, Acrylic acid, esters, polymers
 100-42-5D, Styrene, polymers 106-99-0D, Butadiene, polymers
 123-78-4 302-79-4, all-trans-Retinoic acid 302-79-4D, Retinoic
 acid, derivs. 361-37-5 471-34-1, Calcium carbonate, biological
 studies 1306-06-5, Hydroxylapatite 1332-37-2, Iron oxide,
 biological studies 1404-04-2, Neomycin 4759-48-2, Isotretinoin
 7439-89-6, Iron, biological studies 7439-95-4, Magnesium, biological
 studies 7439-96-5, Manganese, biological studies 7440-25-7,
 Tantalum, biological studies 7440-26-8, Technetium, biological
 studies 7440-39-3, Barium, biological studies 7440-39-3D, Barium,
 compds. 7440-41-7, Beryllium, biological studies 7440-47-3,
 Chromium, biological studies 7440-50-8, Copper, biological studies
 7440-54-2D, Gadolinium, chelates 7631-86-9, Silica, biological
 studies 7778-18-9, Calcium sulfate 9002-72-6, Growth hormone
 9002-86-2, PVC 9003-07-0, Polypropylene 9003-39-8, Plasdone K 90D
 9004-61-9, Hyaluronic acid 9011-14-7, Poly(methyl methacrylate)
~~9012-76-4~~, Chitosan 9061-61-4, NGF 10103-46-5, Calcium
 phosphate 11096-26-7, Erythropoietin 12441-09-7D, Sorbitan, esters
 12619-70-4, Cyclodextrin 14807-96-6, Talc, biological studies
 15802-18-3D, CyanoAcrylic acid, esters, polymers 24980-41-4,
 Polycaprolactone 25248-42-4, Polycaprolactone 25322-68-3,
 Polyethylene glycol 25614-03-3, Bromocriptin 26009-03-0,
 PolyGlycolic acid 26023-30-3,
 Poly[oxy(1-methyl-2-oxo-1,2-ethanediyl)] 26100-51-6, Poly(lactic
 acid) 26124-68-5, PolyGlycolic acid 26354-94-9, Polyvalerolactone

26499-05-8, Polyvalerolactone, SRU 34346-01-5, Glycolic acid-lactic acid copolymer 50903-99-6, L-Name 51110-01-1D, Somatostatin, analogs 59865-13-3, Cyclosporin A 61912-98-9, Insulin-like growth factor 64612-25-5, Fucan 81627-83-0, Macrophage Colony-stimulating factor 83869-56-1, Granulocyte-macrophage Colony-stimulating factor 99896-85-2 114949-22-3, Activin 123626-67-5, Endothelin 1 125265-78-3, N-Carboxybutyl Chitosan 127464-60-2, VEGF 143011-72-7, Granulocyte Colony-stimulating factor 152044-54-7, Epithilone B 154467-38-6 169501-65-9 188492-68-4 189460-40-0, Connective tissue growth factor 250740-90-0, Angiopoietin 302781-03-9 698393-66-7, Isobutylene-styrene triblock copolymer (soft tissue implants and anti-scarring agents)

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L106 ANSWER 3 OF 12 HCA COPYRIGHT 2010 ACS on STN

AN 142:379465 HCA Full-text

TI Prosthetic implants with functionalized carbon surfaces

IN Rathenow, Jorg; Asgari, Soheil; Ban, Andreas; Kunstmann, Jurgen; Mayer, Bernhard

PA Germany

SO U.S. Pat. Appl. Publ., 15 pp., Cont.-in-part of Appl. No. PCT/EP04/05785.

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 10

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 20050079201	A1	20050414	US 2004-939021	20040910
	DE 10324415	A1	20041216	DE 2003-10324415	20030528
	DE 10333098	A1	20050210	DE 2003-10333098	20030721
	DE 10333099	A1	20050210	DE 2003-10333099	20030721
	WO 2004105826	A2	20041209	WO 2004-EP5785	20040528
	WO 2004105826	A3	20050623		
PRAI	DE 2003-10324415	A	20030528		
	DE 2003-10333098	A	20030721		
	DE 2003-10333099	A	20030721		
	WO 2004-EP5785	A2	20040528		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The invention relates to a method of producing medical implants having functionalized surfaces by providing a medical implant with at least one carbon-based layer on at least one part of the surface of the implant, activating the carbon-based layer by creating porosity and functionalizing the activated carbon-based layer. This invention also relates to functionalized implants obtained in by this method (no data).

IT 9012-76-4, Chitosan

(prosthetic implants with functionalized carbon surfaces)

RN 9012-76-4 HCA

CN Chitosan (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

IT 10024-97-2, Nitrous oxide, uses

(prosthetic implants with functionalized carbon surfaces)

RN 10024-97-2 HCA
CN Nitrogen oxide (N2O) (CA INDEX NAME)



IT 9004-34-6, Cellulose, biological studies
(prosthetic implants with functionalized carbon surfaces)

RN 9004-34-6 HCA

CN Cellulose (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

IC ICM B05D003-04

ICS A61F002-02; B05D003-10

INCL 424424000; 623023740; 424426000; 427002210; 427002240

CC 63-7 (Pharmaceuticals)

IT Drug delivery systems

(nanocapsules; prosthetic implants with functionalized
carbon surfaces)

IT Nanostructures

Spheres

(nanospheres; prosthetic implants with functionalized
carbon surfaces)

IT Absorption

Adsorption

Air

Animal cell

Animal tissue

Animal tissue culture

Bone

Cations

Ceramics

Chemisorption

Embryophyta

Emulsions

Ions

Liposomes

Micelles

Microcapsules

Microemulsions

Microorganism

Nanoparticles

Physisorption

Plants

Porosity

Solvents

Sputtering

Viral vectors

(prosthetic implants with functionalized carbon surfaces)

IT 79-41-4D, esters, polymers of 107-73-3, Phosphorylcholine.
7440-02-0, Nickel, biological studies 7440-05-3, Palladium,

biological studies 7440-06-4, Platinum, biological studies
 7440-25-7, Tantalum, biological studies 7440-32-6, Titanium,
 biological studies 7440-44-0, Carbon, biological studies
 7440-48-4, Cobalt, biological studies 7440-50-8, Copper, biological
 studies 7440-57-5, Gold, biological studies 9000-07-1, Carrageenan
 9002-88-4, Polyethylene 9002-89-5 9003-01-4, Polyacrylic acid
 9003-07-0 9004-32-4, Carboxymethyl cellulose 9004-61-9, Hyaluronic
 acid 9004-64-2, Hydroxypropyl cellulose 9004-65-3,
 Hydroxypropylmethyl cellulose 9004-67-5, Methyl cellulose
 9005-25-8, Starch, biological studies 9005-32-7, Alginic acid
~~9012-76-4~~, Chitosan 12597-68-1, Stainless steel, biological
 studies 12683-48-6 24937-78-8, Poly(ethylene vinyl acetate)
 25038-59-9, biological studies 25087-26-7 25104-18-1,
 Poly-L-lysine 25190-06-1, Polytetramethylene glycol 25322-68-3,
 Polyethylene oxide 25322-69-4, Polypropylene oxide 26009-03-0,
 Poly(glycolide) 26023-30-3, Poly[oxy(1-methyl-2-oxo-1,2-ethanediyl)]
 26063-00-3, Poly(hydroxybutyrate) 26202-08-4, Poly(glycolide)
 26680-10-4, Poly(lactide) 26744-04-7 30209-88-2 31621-87-1,
 Polydioxanone 34346-01-5 38000-06-5, Poly-L-lysine 52013-44-2,
 Nitinol 53237-50-6 78644-42-5, Poly(malic acid) 102190-94-3,
 Poly(hydroxyvaleric acid) 111985-13-8 681029-93-6,
 Carboxymethylcellulose phthalate 691397-13-4, Pluronic
 (prosthetic implants with functionalized carbon surfaces)
 IT 1344-28-1, Alumina, uses 7782-44-7, Oxygen, uses ~~10024-97-2~~
 , Nitrous oxide, uses
 (prosthetic implants with functionalized carbon surfaces)
 IT 70-18-8, Glutathione, biological studies 1398-61-4, Chitin
~~9004-34-6~~, Cellulose, biological studies 9004-54-0,
 Dextran, biological studies 9013-20-1, Streptavidin 439211-02-6,
 StrepTactin
 (prosthetic implants with functionalized carbon surfaces)
 OSC.G 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L106 ANSWER 4 OF 12 HCA COPYRIGHT 2010 ACS on STN

AN 142:214882 HCA Full-text

TI Stabilization and ionic triggering of nitric oxide release

IN Smith, Daniel J.

PA The University of Akron, USA

SO PCT Int. Appl., 24 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2005011575	A2	20050210	WO 2004-US23867	20040726
	WO 2005011575	A3	20060112		
	EP 1648527	A2	20060426	EP 2004-779101	20040726
	US 20090136410	A1	20090528	US 2007-565573	20070226
PRAI	US 2003-490218P	P	20030725		
	WO 2004-US23867	W	20040726		


ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Provided is a method for producing nitric oxide that employs an ion exchange resin. Also provided is a method for producing nitric oxide that combines a salt with a gel or cream. A method is provided for producing nitric oxide that combines a pH adjuster with a diazeniumdiolate-contg. compd. or compn.

IT 113-21-3, Lactate, analysis 126-44-3, Citrate, analysis 14265-44-2, Phosphate, analysis
(stabilization and ionic triggering of nitric oxide release)


RN 113-21-3 HCA

CN Propanoic acid, 2-hydroxy-, ion(1-) (CA INDEX NAME)




RN 126-44-3 HCA

CN 1,2,3-Propanetricarboxylic acid, 2-hydroxy-, ion(3-) (CA INDEX NAME)



RN 14265-44-2 HCA

CN Phosphate (CA INDEX NAME)



IT 9000-11-7, CM cellulose 9003-53-6, Polystyrene 9004-34-6, Cellulose, analysis 9012-76-4, Chitosan
(stabilization and ionic triggering of nitric oxide release)

RN 9000-11-7 HCA

CN Cellulose, carboxymethyl ether (CA INDEX NAME)

CM 1

CRN 9004-34-6

CMF Unspecified


CCI PMS, MAN

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

CM 2

CRN 79-14-1

CMF C2 H4 O3




RN 9003-53-6 HCA

CN Benzene, ethenyl-, homopolymer (CA INDEX NAME)

CM 1

CRN 100-42-5

CMF C8 H8



RN 9004-34-6 HCA

CN Cellulose (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

RN 9012-76-4 HCA

CN Chitosan (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

IT ~~10102-43-9~~, Nitric oxide, biological studies

(stabilization and ionic triggering of nitric oxide release)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)




IT ~~839676-39-0~~ ~~839676-40-3~~ ~~839676-41-4~~

(stabilization and ionic triggering of nitric oxide release)


RN 839676-39-0 HCA

CN Propanoic acid, 3-[1-(4-aminobutyl)-2,2-dinitrosohydrazinyl]-, methyl ester, sodium salt (1:1) (CA INDEX NAME)




● Na

RN 839676-40-3 HCA
 CN Propanoic acid, 3-[1-(4-aminobutyl)-2,2-dinitrosohydrazinyl]-, sodium salt (1:1) (CA INDEX NAME)



● Na

RN 839676-41-4 HCA
 CN Propanoic acid, 3-[1-(4-aminobutyl)-2,2-dinitrosohydrazinyl]-, ethyl ester, sodium salt (1:1) (CA INDEX NAME)



● Na

IC ICM A61K
 CC 9-16 (Biochemical Methods)
 IT Ion exchangers
 Nanofibers
 Nanoparticles
 pH
 (stabilization and ionic triggering of nitric oxide release)
 IT 113-21-3, Lactate, analysis 126-44-3, Citrate,
 analysis 14265-44-2, Phosphate, analysis
 (stabilization and ionic triggering of nitric oxide release)
 IT 9000-11-7, CM cellulose 9003-53-6, Polystyrene

9004-34-6, Cellulose, analysis 9012-76-4, Chitosan
 (stabilization and ionic triggering of nitric oxide release)
 IT 10102-43-9, Nitric oxide, biological studies
 (stabilization and ionic triggering of nitric oxide release)
 IT 16545-40-7 27561-78-0 201168-09-4D, Dowex 1X400, reaction with
 NONOates 839676-39-0 839676-40-3
 839676-41-4
 (stabilization and ionic triggering of nitric oxide release)
 OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)
 RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L106 ANSWER 5 OF 12 HCA COPYRIGHT 2010 ACS on STN

AN 142:204864 HCA Full-text
 TI Medical implants coated with porous carbon surfaces carrying drugs
 IN Rathenow, Joerg; Asgari, Soheil; Ban, Andreas
 PA Blue Membranes GmbH, Germany
 SO Ger. Offen., 15 pp.
 CODEN: GWXXBX
 DT Patent
 LA German
 FAN.CNT 10


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	DE 10333099	A1	20050210	DE 2003-10333099	20030721
	DE 202004009061	U1	20040916	DE 2004-202004009061	20040528
	AU 2004243503	A1	20041209	AU 2004-243503	20040528
	CA 2519750	A1	20041209	CA 2004-2519750	20040528
	WO 2004105826	A2	20041209	WO 2004-EP5785	20040528
	WO 2004105826	A3	20050623		
	EP 1626749	A2	20060222	EP 2004-735213	20040528
	EP 1626749	B1	20081008		
	CN 1791436	A	20060621	CN 2004-80013969	20040528
	BR 2004010957	A	20060704	BR 2004-10957	20040528
	JP 2007502184	T	20070208	JP 2006-529943	20040528
	AT 410196	T	20081015	AT 2004-735213	20040528
	PT 1626749	E	20090114	PT 2004-735213	20040528
	EP 2033666	A2	20090311	EP 2008-165943	20040528
	ES 2315661	T3	20090401	ES 2004-735213	20040528
	IL 170898	A	20100328	IL 2004-170898	20040528
	US 20050079201	A1	20050414	US 2004-939021	20040910
	MX 2005011231	A	20060914	MX 2005-11231	20051019
	HK 1089702	A1	20090626	HK 2006-106757	20060613
PRAI	DE 2003-10324415	A1	20030528		
	DE 2003-10333098	A1	20030721		
	DE 2003-10333099	A1	20030721		
	EP 2004-735213	A3	20040528		
	WO 2004-EP5785	W	20040528		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT


AB The invention concerns a method for the prepn. of medical implants with
 functionalized surfaces involving the steps: (a)prepn. of medical implant

that is at least partially coated with a carbon-contg. layer; (b) activation of the carbon-contg. layer by forming a pores on the surface; (c) functionalization of the activated, carbon-contg. surface. The carbon-contg. layer is composed of pyrolytically prepd. carbon, carbon deposited by CVD or PVD process, sputtered carbon, metal carbides, metal carbonitrides, metal oxynitrides, metal oxycarbides or their combinations. The carbon-contg. layers are activated by oxidn. with air, oxygen, dinitrogen oxide, and oxidizing acids, also at elevated temp. A redn. process can also be used for activation. Activated surfaces are functionalized by loading one or more drugs, microorganisms or cells onto the surface. Activated surfaces can be sealed in a CVD or CVI (chem. vapor infiltration) process. The implants are prepd. from carbon, carbon fibers, ceramics, glass, metals, alloys, artificial bone, stone, minerals. Artificial blood vessels, stents, coronary stents, peripheral stents, orthopedic implants, bone and joint prosthesis, artificial heart, heart valves, s.c., and i.m. implants can be activated and functionalized.

IT 10024-97-2, Dinitrogen oxide, biological studies
 (medical implants coated with porous carbon surfaces carrying drugs)
 RN 10024-97-2 HCA
 CN Nitrogen oxide (N2O) (CA INDEX NAME)



IT 9004-34-6, Cellulose, biological studies 9012-76-4, Chitosan 13010-20-3, Nitrosourea
 (medical implants coated with porous carbon surfaces carrying drugs)
 RN 9004-34-6 HCA
 CN Cellulose (CA INDEX NAME)
 *** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
 RN 9012-76-4 HCA
 CN Chitosan (CA INDEX NAME)
 *** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
 RN 13010-20-3 HCA
 CN Urea, N-nitroso- (CA INDEX NAME)



IC ICM A61L027-00
 ICS A61L029-00; A61L033-00; A61F002-30; A61F002-28; A61F002-44;
 A61F002-24
 CC 63-7 (Pharmaceuticals)
 IT Drug delivery systems

(nanocapsules; medical implants coated with porous carbon surfaces carrying drugs)

IT Drug delivery systems

(nanospheres; medical implants coated with porous carbon surfaces carrying drugs)

IT 7782-44-7, Oxygen, biological studies ~~10024-97-2~~, Dinitrogen oxide, biological studies

(medical implants coated with porous carbon surfaces carrying drugs)

IT 50-02-2, Dexamethasone 50-07-7, Mitomycin 50-23-7, Hydrocortisone 50-24-8, Prednisolone 50-56-6, Oxytocin, biological studies 50-78-2, Acetylsalicylic acid 51-41-2, Norepinephrine 51-43-4, Epinephrine 51-45-6, Histamine, biological studies 51-61-6, Dopamine, biological studies 52-53-9, Verapamil 53-03-2, Prednisone 53-06-5, Cortisone 53-86-1, Indomethacin 54-05-7, Chloroquine 56-23-5, Carbon tetrachloride, biological studies 56-54-2, Quinidine 56-75-7, Chloramphenicol 57-22-7, Vincristin 57-41-0, Phenytoine 58-14-0, Pyrimethamin 58-61-7, Adenosine, biological studies 59-05-2, Methotrexate 59-30-3, Folic acid, biological studies 60-54-8, Tetracycline 61-33-6, Penicillin G, biological studies 61-68-7, Mefenamic acid 62-55-5, Thioacetamide 63-74-1, Sulfonamide 64-17-5, Ethanol, biological studies 68-35-9, Sulfadiazine 69-53-4, Ampicillin 71-63-6, Digitoxin 80-08-0, Dapson 83-43-2, Methylprednisolone 87-08-1, Penicillin V 114-07-8, Erythromycin 118-42-3, Hydroxychloroquine 119-04-0, Framycetin 124-94-7, Triamcinolone 127-07-1, Hydroxycarbamide 127-31-1, Fludrocortisone 137-58-6, Lidocaine 140-64-7, Pentamidine diisethionate 152-47-6, Sulfalene 154-21-2, Lincomycin 302-79-4, Tretinoin 356-12-7, Fluocinonide 361-37-5 365-26-4, Oxilofrine 370-14-9, Pholedrine 378-44-9, Betamethasone 382-67-2, Desoximetasone 443-48-1, Metronidazol 466-06-8, Proscillaridin 484-23-1, Dihydralazin 500-92-5, Proguanil 511-12-6, Dihydroergotamine 525-66-6, Propranolol 536-21-0, Norfenefrine 552-94-3, Salsalate 555-30-6, Methyldopa 564-25-0, Doxycycline 586-06-1, Orciprenaline 630-60-4, Ouabain 638-94-8, Desonide 644-62-2 660-27-5, Diisopropyl amine dichloroacetate 709-55-7, Etilefrine 738-70-5, Trimethoprim 768-94-5, Amantadine 807-38-5, Fluocinolone 865-21-4, Vinblastin 1066-17-7, Colistin 1306-05-4, Fluorapatite 1306-06-5, Hydroxylapatite 1393-87-9, Fusafungine 1404-26-8, Polymyxin B 1404-90-6, Vancomycin 1524-88-5, Flurandrenolide 1695-77-8, Spectinomycin 1951-25-3, Amiodarone 2589-47-1, Prajmaliumbitartrate, biological studies 2809-21-4, Etidronic acid 3056-17-5, Stavudine 3093-35-4, Halcinonide 3385-03-3, Flunisolide 3737-09-5, Disopyramide 3930-20-9, Sotalol 4360-12-7, Ajmalin 4419-39-0, Beclomethasone 4828-27-7, Clo cortolone 4936-47-4, Nifuratel 5104-49-4, Flurbiprofen 5355-48-6 6452-71-7, Oxprenolol 6990-06-3, Fusidinic acid 7440-22-4, Silver, biological studies 7440-25-7, Tantalum, biological studies 7440-32-6, Titanium, biological studies 7440-66-6, Zinc, biological studies 7481-89-2, Zalcitabine 7542-37-2, Paromomycin 7681-49-4, Sodium fluoride, biological studies 7758-87-4, Tricalciumphosphate 8001-27-2, Hirudin

8025-81-8, Spiramycin 8067-24-1, Co-Dergocrine mesylate 9000-07-1, Carrageenan 9002-01-1, Streptokinase 9002-60-2, Corticotropin, biological studies 9002-71-5, Thyrotrophin 9002-88-4, Polyethylene 9002-89-5, Polyvinylalcohol 9003-01-4, Acrylic acid homopolymer 9003-07-0, Polypropylene 9003-39-8, Polyvinylpyrrolidone 9004-32-4, Carboxymethylcellulose ~~9004-34-6~~, Cellulose, biological studies 9004-54-0, Dextran, biological studies 9004-61-9, Hyaluronic acid 9004-64-2, Hydroxypropylcellulose 9004-65-3, Hydroxypropylmethylcellulose 9004-67-5, Methylcellulose 9005-25-8, Starch, biological studies 9005-32-7, Alginic acid 9005-49-6, Heparin, biological studies ~~9012-76-4~~, Chitosan 9039-53-6, Urokinase 9061-61-4, Nerve growth factor 10118-90-8, Minocycline 10163-15-2, Disodium fluorophosphate 10596-23-3, Clodronic acid 11096-26-7, Erythropoietin 11111-12-9, Cephalosporin 11128-99-7, Angiotensin II 12597-68-1, Stainless steel, biological studies 12629-01-5, Somatropin 12683-48-6 ~~13010-20-3~~, Nitrosourea 13292-46-1, Rifampicin 13463-67-7, Titanium dioxide, biological studies 14402-89-2, Nitroprusside sodium 14636-12-5, Terlipressin 15307-86-5, Diclofenac 15663-27-1, Cisplatin 15686-71-2, Cefalexin 15687-27-1, Ibuprofen 16662-47-8, Gallopamil 16679-58-6, Desmopressin 16846-24-5, Josamycin 18323-44-9, Clindamycin 19216-56-9, Prazosin 19387-91-8, Tinidazol 19388-87-5, Taurolidine 20830-75-5, Digoxin 21256-18-8, Oxaprozin 21829-25-4, Nifedipine 22071-15-4, Ketoprofen 22204-53-1, Naproxen 22254-24-6, Ipratropium bromide 22494-42-4, Diflunisal 23155-02-4, Fosfomycin 24937-78-8 25038-59-9, biological studies 25087-26-7, Methacrylic acid homopolymer 25104-18-1, Polylysine 25122-41-2, Clobetasol 25190-06-1, Poly(Tetramethylene glycol) 25322-68-3, Polyethylene oxide 25322-69-4, Polypropylene oxide 25614-03-3, Bromocriptine 25953-19-9, Cefazolin 26009-03-0, Polyglycolic acid 26023-30-3, Poly[oxy(1-methyl-2-oxo-1,2-ethanediyl)] 26063-00-3, β -Hydroxybutyric acid homopolymer 26099-09-2 26100-51-6, Polylactic acid 26124-68-5, Polyglycolic acid 26171-23-3, Tolmetin 26744-04-7, β -Hydroxybutyric acid homopolymer, sru 26787-78-0, Amoxicillin 26807-65-8, Indapamide 26844-12-2, Indoramin 29122-68-7, Atenolol 29679-58-1, Fenoprofen 30209-88-2, Polyallyl cyanoacrylate 30516-87-1, Zidovudine 30578-37-1, Amezinium metil sulfate 30685-43-9, Metildigoxin 31621-87-1, Polydioxanone 31828-71-4, Mexiletine 33069-62-4, Paclitaxel 33515-09-2, Gonadorelin 33774-52-6, Detajmumbitartrate, biological studies 34346-01-5, Lactic acid-glycolic acid copolymer 34661-75-1, Urapidil 35607-66-0, Cefoxitin 36322-90-4, Piroxicam 36703-88-5 36791-04-5, Ribavirin 38194-50-2, Sulindac 38304-91-5, Minoxidil 39562-70-4, Nitrendipine 40391-99-9 41340-25-4, Etodolac 41575-94-4, Carboplatin 41708-72-9, Tocainide 42399-41-7, Diltiazem 42794-76-3, Midodrine 42924-53-8, Nabumetone 50370-12-2, Cefadroxil 50972-17-3, Bacampicillin 51022-69-6, Amcinonide 51110-01-1, Somatostatin 51264-14-3, Amsacrine 51333-22-3, Budesonide 51384-51-1, Metoprolol 51481-65-3, Mezlocillin 51940-44-4, Pipemidic acid 52013-44-2, Nitinol

53123-88-9, Sirolimus 53230-10-7, Mefloquine 53237-50-6
 53714-56-0, Leuprorelin 53910-25-1, Pentostatin 53994-73-3,
 Cefaclor 54063-53-5, Propafenone 54143-55-4, Flecainide
 54143-56-5, Flecainide acetate 55142-85-3, Ticlopidine 55268-75-2,
 Cefuroxim 56391-56-1, Netilmicin 57773-63-4, Triptorelin
 57982-77-1, Buserelin 58066-85-6, Miltefosine 59277-89-3,
 Aciclovir 61036-62-2, Teicoplanin 61477-96-1, Piperacillin
 61622-34-2, Cefotiam

(medical implants coated with porous carbon surfaces carrying
 drugs)

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L106 ANSWER 6 OF 12 HCA COPYRIGHT 2010 ACS on STN

AN 142:42564 HCA Full-text

TI Treatment of carbon **nanosstructure** using fluidization

IN Jung, Kyeong Taek; Kim, Myung Soo; Jeon, Kwan Goo; Lee, Young Hee

PA S. Korea

SO U.S. Pat. Appl. Publ., 13 pp.

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 20040253374	A1	20041216	US 2004-830914	20040423
	KR 2004091951	A	20041103	KR 2003-25733	20030423
	KR 2004093542	A	20041106	KR 2003-27453	20030430
	JP 2005001980	A	20050106	JP 2004-128506	20040423
PRAI	KR 2003-25733	A	20030423		
	KR 2003-27453	A	20030430		

AB The present invention relates to an efficient and simple method for treating
 a carbon **nanosstructure** by fluidizing the carbon **nanosstructure** in a reactor
 using a carrier gas and a reactive gas to contact the fluidized carbon
nanosstructure. Carbon **nanosstructures** can be effectively purified, uniformly
 surface-treated and easily employable in the post-process, e.g., in the
 prodn. of a composite.

IT ~~10024-97-2~~, Nitrogen oxide (N2O), processes ~~10102-43-9~~
 , Nitrogen oxide (NO), processes ~~10102-44-0~~, Nitrogen oxide
 (NO2), processes
 (surface treating agent; treatment of carbon **nanosstructure**
 using fluidization)

RN 10024-97-2 HCA

CN Nitrogen oxide (N2O) (CA INDEX NAME)


○══N══N

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



RN 10102-44-0 HCA
 CN Nitrogen oxide (NO2) (CA INDEX NAME)



IT 9000-11-7, Carboxymethyl cellulose
 (treatment of carbon ~~nanos~~tructure using fluidization)
 RN 9000-11-7 HCA
 CN Cellulose, carboxymethyl ether (CA INDEX NAME)


CM 1

CRN 9004-34-6
 CMF Unspecified
 CCI PMS, MAN

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

CM 2

CRN 79-14-1
 CMF C2 H4 O3



IC ICM C23C016-26
 INCL 427213000; 427249100
 CC 57-8 (Ceramics)
 Section cross-reference(s): 66
 ST carbon ~~nanos~~tructure fluidization surface treatment
 composite manuf
 IT Sulfonation
 (agent; treatment of carbon ~~nanos~~tructure using
 fluidization)
 IT Titanates
 (alkoxides, secondary surface treatment agent; treatment of carbon
~~nanos~~tructure using fluidization)
 IT Silanes
 (alkoxy, secondary surface treatment agent; treatment of carbon
~~nanos~~tructure using fluidization)

IT Metal alkoxides
 (aluminum, secondary surface treatment agent; treatment of carbon
 nanostructure using fluidization)

IT Nanostructures
 (carbon; treatment of carbon nanostructure using
 fluidization)

IT Gases
 (carrier; treatment of carbon nanostructure using
 fluidization)

IT Vapor deposition process
 (chem.; treatment of carbon nanostructure using
 fluidization)

IT Air
 (purifying gas; treatment of carbon nanostructure using
 fluidization)

IT Composites
 (reinforced; treatment of carbon nanostructure using
 fluidization)

IT Carbonates, processes
 Chlorides, processes
 Metal alkoxides
 Nitrates, processes
 Phosphines
 (secondary surface treatment agent; treatment of carbon
 nanostructure using fluidization)

IT Metal alkoxides
 (titanium, secondary surface treatment agent; treatment of carbon
 nanostructure using fluidization)

IT Coupling agents
 Dispersion (of materials)
 Etching
 Fluidization
 Fluidized beds
 Fluorination
 Heat treatment
 Nitration
 Oxidation
 Plasma
 Purification
 Raman spectra
 Surface treatment
 X-ray photoelectron spectra
 (treatment of carbon nanostructure using fluidization)

IT Metals, processes
 (vaporized, secondary surface treatment agent; treatment of carbon
 nanostructure using fluidization)

IT 1333-74-0, Hydrogen, processes 7664-41-7, Ammonia, processes
 (etching gas; treatment of carbon nanostructure using
 fluidization)

IT 7440-37-1, Argon, uses 7440-59-7, Helium, uses 7727-37-9,
 Nitrogen, uses
 (gas carrier; treatment of carbon nanostructure using

fluidization)

IT 7440-44-0P, Carbon, preparation
(**nanost**tructure; treatment of carbon **nanost**tructure
using fluidization)

IT 124-38-9, Carbon dioxide, processes 7647-01-0, Hydrochloric acid,
processes 7664-39-3, Fluorhydric acid, processes 7664-93-9,
Sulfuric acid, processes 7697-37-2, Nitric acid, processes
7722-84-1, Hydrogen peroxide, processes
(purifying gas and surface treating agent; treatment of carbon
nanosttructure using fluidization)

IT 7782-44-7, Oxygen, processes
(purifying gas; treatment of carbon **nanost**tructure using
fluidization)

IT 71-50-1, Acetate, processes
(secondary surface treatment agent; treatment of carbon
nanosttructure using fluidization)

IT 74-90-8, Hydrogen cyanide, processes 110-86-1, Pyridine, processes
7446-09-5, Sulfur oxide, processes 7664-38-2, Phosphoric acid,
processes 7722-64-7, Potassium permanganate 7758-05-6, Potassium
iodate 7782-50-5, Chlorine, processes 7783-06-4, Hydrogen sulfide,
processes 10024-97-2, Nitrogen oxide (N2O), processes
10028-15-6, Ozone, processes 10049-04-4, Chlorine dioxide
10102-43-9, Nitrogen oxide (NO), processes 10102-44-0
, Nitrogen oxide (NO2), processes 12624-32-7, Sulfur oxide
(surface treating agent; treatment of carbon **nanost**tructure
using fluidization)

IT 102-54-5, Ferrocene
(treatment of carbon **nanost**tructure using fluidization)

IT 64-17-5, Ethanol, processes 71-43-2, Benzene, processes 78-10-4,
TEOS 7782-41-4, Fluorine, processes 10026-04-7, Tetrachlorosilane
(treatment of carbon **nanost**tructure using fluidization)

IT 9000-11-7, Carboxymethyl cellulose 25155-30-0, Sodium
dodecyl-benzene sulfonate
(treatment of carbon **nanost**tructure using fluidization)

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L106 ANSWER 7 OF 12 HCA COPYRIGHT 2010 ACS on STN

AN 141:413669 HCA Full-text

TI Fuel cell component with lyophilic surface

IN Extrand, Charles W.; Bhatt, Sanjiv M.; Monson, Loxie

PA Entegris, Inc., USA

SO PCT Int. Appl., 33 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2004100287	A2	20041118	WO 2004-US13560	20040503
	WO 2004100287	A3	20050526		
	US 20040258975	A1	20041223	US 2004-837241	20040430
PRAI	US 2003-468213P	P	20030505		

US 2004-837241 A 20040430

AB A fuel cell component with surfaces having improved lyophilicity is disclosed so that liq. on the component adheres closely to the surface in relatively flat droplets or sheets. The lyophilic surfaces may be formed by cold plasma or UV light treatment of the component. The lyophilic surfaces may be selectively provided on crit. areas of the component, such as for example on flow channel wall surfaces of bipolar plates and membrane electrode assemblies, thereby inhibiting liq. blocking of the flow channels during operation of the fuel cell.

IT 9003-53-6, Polystyrene
(fuel cell component with lyophilic surface)


RN 9003-53-6 HCA

CN Benzene, ethenyl-, homopolymer (CA INDEX NAME)

CM 1

CRN 100-42-5


CMF C8 H8



IT 10024-97-2, Nitrous oxide, uses
(process gas; fuel cell component with lyophilic surface)

RN 10024-97-2 HCA

CN Nitrogen oxide (N2O) (CA INDEX NAME)



IC ICM H01M

CC 52-2 (Electrochemical, Radiational, and Thermal Energy Technology)
Section cross-reference(s): 38

IT **Nanotubes**

(carbon, filler; fuel cell component with lyophilic surface)

IT 57-13-6, Urea, uses 100-42-5D, Styrene, block copolymers, with
olefins 126-99-8, Chloroprene 131-17-9, Diallyl phthalate
9002-84-0, Polytetrafluoroethylene 9002-86-2, Polyvinyl chloride
9003-17-2, Polybutadiene 9003-31-0, Polyisoprene 9003-53-6
, Polystyrene 24937-79-9, PvdF 25778-04-5 413569-08-1, uses
(fuel cell component with lyophilic surface)

IT 7440-44-0, Carbon, uses
(nanotubes, filler; fuel cell component with lyophilic
surface)

IT 56-23-5, Carbon tetrachloride, uses 75-21-8, Ethylene oxide, uses
107-18-6, Allyl alcohol, uses 107-21-1, Ethylene glycol, uses
115-10-6, Methyl ether 124-38-9, Carbon dioxide, uses 630-08-0,
Carbon monoxide, uses 7440-37-1, Argon, uses 7446-09-5, Sulfur

oxide, uses 7664-41-7, Ammonia, uses 7727-37-9, Nitrogen, uses 7782-44-7, Oxygen, uses 7782-50-5, Chlorine, uses 10024-97-2, Nitrous oxide, uses 10028-15-6, Ozone, uses 10049-04-4, Chlorine dioxide 11104-93-1, Nitrogen oxide, uses

(process gas; fuel cell component with lyophilic surface)

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L106 ANSWER 8 OF 12 HCA COPYRIGHT 2010 ACS on STN

AN 141:266048 HCA Full-text

TI Medical implants with carbon-containing surfaces that are functionalized

PA Blue Membranes GmbH, Germany

SO Ger. Gebrauchsmusterschrift, 18 pp.

CODEN: GGXXFR

DT Patent

LA German

FAN.CNT 10

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	DE 202004009061	U1	20040916	DE 2004-202004009061	20040528
	DE 10324415	A1	20041216	DE 2003-10324415	20030528
	DE 10333098	A1	20050210	DE 2003-10333098	20030721
	DE 10333099	A1	20050210	DE 2003-10333099	20030721
PRAI	DE 2003-10324415	A1	20030528		
	DE 2003-10333098	A1	20030721		
	DE 2003-10333099	A1	20030721		

AB The invention concerns medical implants with carbon-contg. surfaces that are functionalized; the surfaces are prepd. by (a) prepg. a medical implant with a carbon-contg. surface; (b) activation of the carbon layer by creating porosity; (c) functionalization of the activated, carbon-contg. layer. The carbon layer can be prepd. by pyrolysis, CVD, PVD, sputtering, ion implantation. The medical devices are prepd. from carbon, carbon-composite material, glass, ceramics, glass fibers, carbon fibers, metals, stainless steel, titanium, tantalum, platinum, nitinol, alloys, artificial bone, minerals, and their combinations. Artificial blood vessels, stents, coronary stents, peripheral stents, orthopedic implants, artificial hearts and heart valves, artificial bones and joints are prepd. The carbon layer is activated with oxidn. or reducing agents in the presence of air, oxygen, nitrogen monoxide, oxidative acids; heat and/or ultrasound can be applied. The activated implant surfaces are functionalized with drugs, microorganisms, plant, animal or human cells. The invention also concerns controlled-release implanted drug delivery systems.


IT 10102-43-9, Nitrogen monoxide, biological studies

(medical implants with carbon-contg. surfaces that are functionalized)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

IT 9004-34-6D, Cellulose, derivs. 9012-76-4, Chitosan
 13010-20-3, Nitrosourea
 (medical implants with carbon-contg. surfaces that are
 functionalized)
 RN 9004-34-6 HCA
 CN Cellulose (CA INDEX NAME)
 *** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
 RN 9012-76-4 HCA
 CN Chitosan (CA INDEX NAME)
 *** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
 RN 13010-20-3 HCA
 CN Urea, N-nitroso- (CA INDEX NAME)



IC ICM A61L027-50
 CC 63-7 (Pharmaceuticals)
 IT Drug delivery systems
 (nanocapsules; medical implants with carbon-contg.
 surfaces that are functionalized)
 IT 7440-21-3, Silicon, biological studies 7440-44-0, Carbon, biological
 studies 7782-44-7, Oxygen, biological studies 10102-43-9,
 Nitrogen monoxide, biological studies
 (medical implants with carbon-contg. surfaces that are
 functionalized)
 IT 50-02-2, Dexamethasone 50-23-7, Hydrocortisone 50-24-8,
 Prednisolone 50-56-6, Oxytocin, biological studies 50-78-2,
 Acetylsalicylic acid 51-41-2, Norepinephrine 51-43-4, Epinephrine
 51-45-6, Histamine, biological studies 51-61-6, Dopamin, biological
 studies 52-53-9, Verapamil 53-03-2, Prednisone 53-06-5,
 Cortisone 53-86-1, Indomethacin 54-05-7, Chloroquine 56-23-5,
 Carbon tetrachloride, biological studies 56-54-2, Quinidine
 56-75-7, Chloramphenicol 57-22-7, Vincristin 57-41-0, Phenytoin
 57-62-5 57-92-1, Streptomycin, biological studies 58-14-0,
 Pyrimethamine 58-61-7, Adenosine, biological studies 59-05-2,
 Methotrexate 59-30-3, Folic acid, biological studies 60-54-8,
 Tetracycline 60-54-8D, Tetracycline, derivs. 61-33-6, Penicillin
 G, biological studies 61-68-7, Mefenamic acid 62-55-5,
 Thioacetamide 63-74-1, Sulfonamide 64-17-5, Ethanol, biological
 studies 67-96-9, Dihydrotachysterol 68-35-9, Sulfadiazine
 69-53-4, Ampicillin 70-18-8, Glutathione, biological studies
 71-63-6, Digitoxin 79-10-7D, Acrylic acid, esters, polymers
 79-41-4D, Methacrylic acid, esters, polymers 79-57-2,
 Oxytetracycline 80-08-0, Dapson 83-43-2, Methylprednisolone
 87-08-1, Penicillin V 108-05-4D, Vinylacetate, copolymers with

phthalates 114-07-8, Erythromycin 118-42-3, Hydroxychloroquine
 119-04-0, Framycetin 120-73-0D, Purine, derivs. 124-94-7,
 Triamcinolone 127-07-1, Hydroxycarbamide 127-31-1, Fludrocortisone
 130-95-0D, Quinine, derivs. 137-58-6, Lidocaine 140-64-7,
 Pentamidine diisethionate 154-21-2, Lincomycin 289-95-2D,
 Pyrimidine, derivs. 302-79-4, Tretinoin 356-12-7, Fluocinonide
 361-37-5 365-26-4, Oxilofrine 370-14-9, Pholedrine 378-44-9,
 Betamethasone 382-67-2, Desoximetasone 443-48-1, Metronidazol
 466-06-8 484-23-1, Dihydralazin 500-92-5, Proguanil 511-12-6,
 Dihydroergotamine 525-66-6, Propranolol 536-21-0, Norfenefrine
 552-94-3, Salsalate 555-30-6, Methyldopa 564-25-0, Doxycycline
 586-06-1, Orciprenaline 630-60-4, Ouabain 638-94-8, Desonide
 644-62-2 660-27-5, Diisopropyl amine dichloroacetate 709-55-7,
 Etilefrine 738-70-5, Trimethoprim 768-94-5, Amantadine 807-38-5,
 Fluocinolone 865-21-4, Vinblastin 1066-17-7, Colistin 1344-28-1,
 Alumina, biological studies 1393-87-9, Fusafungin 1403-66-3,
 Gentamicin 1404-00-8, Mitomycin 1404-04-2, Neomycin 1404-26-8,
 Polymyxin-B 1404-90-6, Vancomycin 1406-05-9, Penicillin
 1524-88-5, Flurandrenolide 1695-77-8, Spectinomycin 1951-25-3,
 Amiodarone 2589-47-1, Prajmaliumbitartrate, biological studies
 2809-21-4, Etidronic acid 3056-17-5, Stavudine 3093-35-4,
 Halcinonide 3385-03-3, Flunisolid 3737-09-5, Disopyramide
 3930-20-9, Sotalol 4360-12-7, Ajmalin 4419-39-0, Beclomethasone
 4428-95-9, Foscarnet 4828-27-7, Clocortolone 4936-47-4, Nifuratel
 5104-49-4, Flurbiprofen 5355-48-6 6452-71-7, Oxprenolol
 6990-06-3, Fusidinic acid 7440-02-0, Nickel, biological studies
 7440-06-4, Platinum, biological studies 7440-22-4, Silver,
 biological studies 7440-25-7, Tantalum, biological studies
 7440-32-6, Titanium, biological studies 7440-41-7, Beryllium,
 biological studies 7440-48-4, Cobalt, biological studies
 7440-50-8, Copper, biological studies 7481-89-2, Zalcitabine
 7542-37-2, Paromomycin 7631-86-9, Silica, biological studies
 7681-49-4, Sodium fluoride, biological studies 8001-27-2, Hirudin
 8025-81-8, Spiramycin 8067-24-1, Dihydroergotoxine methane sulfonate
 9000-94-6, Antithrombin 9001-90-5, Plasmin 9002-01-1,
 Streptokinase 9002-60-2, Corticotropin, biological studies
 9002-71-5, Thyrotrophin 9002-72-6, Growth hormone 9002-88-4,
 Polyethylene 9002-89-5, Polyvinylalcohol 9003-07-0, Polypropylene
 9003-28-5, Polybutene 9003-39-8, Polyvinylpyrrolidone
~~9004-34-6D~~, Cellulose, derivs. 9004-54-0, Dextran,
 biological studies 9004-61-9, Hyaluronic acid 9004-64-2,
 Hydroxypropylcellulose 9004-65-3, Hydroxypropylmethylcellulose
 9004-67-5, Methylcellulose 9005-25-8, Starch, biological studies
 9005-49-6, Heparin, biological studies 9007-12-9, Calcitonin
~~9012-76-4~~, Chitosan 9013-20-1, Streptavidin 9039-53-6,
 Urokinase 9061-61-4, NGF 10118-90-8, Minocycline 10163-15-2,
 Disodium fluorophosphate 10596-23-3, Clodronic acid 11056-06-7,
 Bleomycin 11096-26-7, Erythropoietin 11111-12-9, Cephalosporin
 11128-99-7, Angiotensin II 12597-68-1, Stainless steel, biological
 studies 12629-01-5, Somatropin 12683-48-6 ~~13010-20-3~~,
 Nitrosourea 13292-46-1, Rifampicin 13463-67-7, Titanium dioxide,
 biological studies 14402-89-2, Nitroprusside sodium 14636-12-5,

Terlipressin 15307-86-5, Diclofenac 15663-27-1, Cisplatin 15686-71-2, Cefalexin 15687-27-1, Ibuprofen 15802-18-3, 16662-47-8, Gallopamil 16679-58-6, Desmopressin 16846-24-5, Josamycin 18323-44-9, Clindamycin 19216-56-9, Prazosin 19387-91-8, Tinidazol 19388-87-5, Taurolidine 20830-75-5, Digoxin 20830-81-3, Daunorubicin 21256-18-8, Oxaprozin 21829-25-4, Nifedipine 22071-15-4, Ketoprofen 22204-53-1, Naproxen 22254-24-6, Ipratropium bromide 22494-42-4, Diflunisal 23155-02-4, Fosfomycin 23214-92-8, Doxorubicin 24937-78-8, Polyethylenevinyl acetate 25014-41-9, 2-Propenenitrile, homopolymer 25038-59-9, Polyethyleneterephthalate, biological studies 25122-41-2, Clobetasol 25190-06-1, Polytetramethylene glycol 25322-68-3, Polyethylene oxide 25322-69-4, Polypropylene oxide 25614-03-3, Bromocriptine 25953-19-9, Cefazolin 26009-03-0, Polyglycolide 26023-30-3, D,L-Lactic acid, homopolymer 26063-00-3, Polyhydroxybutyrate 26099-09-2, Polymaleic acid 26100-51-6, Polylactic acid 26171-23-3, Tolmetin 26202-08-4, Polyglycolide 26744-04-7, 26787-78-0, Amoxicillin 26807-65-8, Indapamide 26844-12-2, Indoramin 29122-68-7, Atenolol 29679-58-1, Fenoprofen 30209-88-2, 30516-87-1, Zidovudine 30578-37-1, Amezinium methyl sulfate 30685-43-9, Metildigoxin 31621-87-1, Polydioxanone 31828-71-4, Mexiletine 32986-56-4, Tobramycin 33069-62-4, Paclitaxel 33515-09-2, Gonadorelin 33774-52-6, Detajmiumbitartrate, biological studies 34346-01-5, Glycolic acid-lactic acid copolymer 34368-04-2, Dobutamine 34661-75-1, Urapidil 35607-66-0, Cefoxitin 36322-90-4, Piroxicam 36703-88-5, 36791-04-5, Ribavirin 36877-68-6D, Nitroimidazole, derivs. 37203-62-6, Blood coagulation factor XIIa 37517-28-5, Amikacin 38000-06-5, Polylysine 38194-50-2, Sulindac 38304-91-5, Minoxidil 39562-70-4, Nitrendipine 40391-99-9, 41340-25-4, Etodolac 41575-94-4, Carboplatin 42399-41-7, Diltiazem 42794-76-3, Midodrine 42924-53-8, Nabumetone 50370-12-2, Cefadroxil (medical implants with carbon-contg. surfaces that are functionalized)

L106 ANSWER 9 OF 12 HCA COPYRIGHT 2010 ACS on STN

AN 140:79159 HCA [Full-text](#)

TI Particles from supercritical fluid extraction of emulsion

IN Chattopadhyay, Pratibhash; Shekunov, Boris Y.; Seitzinger, Jeffrey S.; Huff, Robert W.

PA Ferro Corporation, USA

SO PCT Int. Appl., 61 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 4

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2004004862	A1	20040115	WO 2003-US19633	20030620
	US 20040026319	A1	20040212	US 2003-423492	20030425
	US 6998051	B2	20060214		
	CA 2483563	A1	20040115	CA 2003-2483563	20030620

CA 2483563	C	20080826		
AU 2003281210	A1	20040123	AU 2003-281210	20030620
EP 1551523	A1	20050713	EP 2003-742125	20030620
EP 1551523	B1	20070808		
CN 1665576	A	20050907	CN 2003-815675	20030620
CN 1318116	C	20070530		
JP 2005531408	T	20051020	JP 2004-519622	20030620
JP 4421475	B2	20100224		
AT 369198	T	20070815	AT 2003-742125	20030620
ES 2289308	T3	20080201	ES 2003-742125	20030620
PRAI US 2002-393904P	P	20020703		
US 2003-445944P	P	20030207		
US 2003-423492	A	20030425		
WO 2003-US19633	W	20030620		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A method of producing microparticles and ~~nanoparticles~~ of a solute via the extn. of solvent, having the solute dissolved therein, from an emulsion fed to a vessel using a supercrit. fluid also fed to the vessel. The solute to be pptd. is dissolved in the solvent to form a soln., and the soln. is dispersed in an immiscible or partially miscible liq. to form an emulsion which is fed by a tube to the vessel. The particles are produced via the extn. of the solvent from the emulsion using the supercrit. fluid in the vessel. The process can produce an aq. suspension of particles that are substantially insol. in water, and the solvents used in the process to form the emulsion initially can be recovered and recycled from vessel ports at the top.

IT 9003-53-6, Polystyrene
(~~nanoparticle~~ formation of; ~~nanoparticles~~ from supercrit. fluid extn. of emulsion)


RN 9003-53-6 HCA

CN Benzene, ethenyl-, homopolymer (CA INDEX NAME)

CM 1

CRN 100-42-5


CMF C8 H8



IT 10024-97-2, Nitrous oxide, processes
(particles from supercrit. from supercrit. fluid extn. of emulsion)

RN 10024-97-2 HCA

CN Nitrogen oxide (N2O) (CA INDEX NAME)



IC ICM B01D011-04
 CC 48-6 (Unit Operations and Processes)
 Section cross-reference(s): 17, 38, 45, 50, 64, 66
 ST particle supercrit fluid emulsion extn **nanoparticle** CO2
 solvent colloid
 IT Natural products, pharmaceutical
 (**nanoparticle** formation of animal or plant exts.;
 particles from supercrit. from supercrit. fluid extn. of emulsion)
 IT Polymers, processes
 (**nanoparticle** formation of precursors; particles from
 supercrit. from supercrit. fluid extn. of emulsion)
 IT Virus
 (**nanoparticle** formation of viral materials; particles
 from supercrit. from supercrit. fluid extn. of emulsion)
 IT Agrochemicals
 Antibiotics
 Biodegradable materials
 Catalysts
 Cosmetics
 Diagnostic agents
 Dietary supplements
 Drugs
 Dyes
 Explosives
 Insecticides
 Paints
 Pigments, nonbiological
 (**nanoparticle** formation of; particles from supercrit.
 from supercrit. fluid extn. of emulsion)
 IT Alkaloids, processes
 Antigens
 Enzymes, processes
 Lipids, processes
 Nucleic acids
 Peptides, processes
 Polymers, processes
 Proteins
 Toxins
 Vitamins
 (**nanoparticle** formation of; particles from supercrit.
 from supercrit. fluid extn. of emulsion)
 IT Emulsions
 Nanoparticles
 Precipitation (chemical)
 Supercritical fluids
 Tanks (containers)
 (**nanoparticles** from supercrit. fluid extn. of emulsion)
 IT Drug delivery systems
 (**nanoparticles**, **nanoparticle** formation of;
 particles from supercrit. from supercrit. fluid extn. of emulsion)
 IT **Nanostructures**
 Spheres

(nanospheres; particles from supercrit. from supercrit. fluid extn. of emulsion)

IT Solvents
(non-polar and partially water sol.; nanoparticles from supercrit. fluid extn. of emulsion)

IT Extraction
(supercrit.; nanoparticles from supercrit. fluid extn. of emulsion)

IT 9003-53-6, Polystyrene
(nanoparticle formation of; nanoparticles from supercrit. fluid extn. of emulsion)

IT 555-44-2, Tripalmitin 604-35-3, Cholesterol Acetate 33434-24-1, EUDRAGIT RS 34346-01-5, Glycolic Acid-Lactic acid copolymer
(nanoparticle formation of; particles from supercrit. from supercrit. fluid extn. of emulsion)

IT 67-66-3, Chloroform, processes 75-46-7, Trifluoromethane 108-88-3, Toluene, processes 115-10-6, Dimethyl ether 141-78-6, Ethyl Acetate, processes 10024-97-2, Nitrous oxide, processes
(particles from supercrit. from supercrit. fluid extn. of emulsion)

OSC.G 12 THERE ARE 12 CAPLUS RECORDS THAT CITE THIS RECORD (14 CITINGS)

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L106 ANSWER 10 OF 12 HCA COPYRIGHT 2010 ACS on STN

AN 133:122599 HCA Full-text

TI Carbide and oxycarbide based compositions and nanorods

IN Moy, David; Niu, Chun-Ming; Ma, Jun; Willey, Jason M.

PA Hyperion Catalysis International, Inc., USA

SO PCT Int. Appl., 69 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 5

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2000041808	A1	20000720	WO 2000-US753	20000112
	CA 2359336	A1	20000720	CA 2000-2359336	20000112
	EP 1152827	A1	20011114	EP 2000-903266	20000112
	JP 2002534351	T	20021015	JP 2000-593411	20000112
	AU 764311	B2	20030814	AU 2000-25040	20000112
	EP 1920837	A2	20080514	EP 2007-122314	20000112
	EP 1920837	A3	20081119		
	KR 907214	B1	20090710	KR 2001-708727	20000112
	MX 2001007030	A	20020311	MX 2001-7030	20010711
PRAI	US 1999-115735P	P	19990112		
	EP 2000-903266	A3	20000112		
	WO 2000-US753	W	20000112		

AB Compns. including oxycarbide-based nanorods and/or carbide-based nanorods and/or carbon nanotubes bearing carbides and oxycarbides and methods of making the same are provided. Rigid porous structures including oxycarbide-based nanorods and/or carbide based nanorods and/or carbon nanotubes bearing

carbides and oxycarbides and methods of making the same are also provided. The compns. and rigid porous structures of the invention can be used either as catalyst and/or catalyst supports in fluid phase catalytic chem. reactions. Processes for making supported catalyst for selected fluid phase catalytic reactions are also provided. The fluid phase catalytic reactions catalyzed include hydrogenation, hydrodesulfurization, hydrodenitrogenation, hydrodemetallization, hydrodeoxygenation, hydrodearomatization, dehydrogenation, hydrogenolysis, isomerization, alkylation, dealkylation and transalkylation.

IT 9003-53-6, Polystyrene 9004-34-6, Cellulose,
reactions
(carbide and oxycarbide based compns. and nanorods)


RN 9003-53-6 HCA

CN Benzene, ethenyl-, homopolymer (CA INDEX NAME)

CM 1

CRN 100-42-5

CMF C8 H8



RN 9004-34-6 HCA


CN Cellulose (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

IT 10024-97-2, Nitrous oxide, uses 10102-43-9, Nitric
oxide, uses 10102-44-0, Nitrogen dioxide, uses
(oxidant; carbide and oxycarbide based compns. and nanorods
)

RN 10024-97-2 HCA

CN Nitrogen oxide (N2O) (CA INDEX NAME)



RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



RN 10102-44-0 HCA

CN Nitrogen oxide (NO2) (CA INDEX NAME)

O—N≡O

IC ICM B01J027-22
ICS C01B013-14; C03B025-00

CC 51-4 (Fossil Fuels, Derivatives, and Related Products)
Section cross-reference(s): 67

ST carbide oxycarbide ~~nanorod~~ catalyst

IT Alkylation catalysts
Dealkylation catalysts
Dehydrogenation catalysts
Hydrogenation catalysts
Isomerization catalysts
Transalkylation catalysts
(carbide and oxycarbide based compns. and ~~nanorods~~)

IT Carbides
(carbide and oxycarbide based compns. and ~~nanorods~~)

IT Carbohydrates, reactions
Phenolic resins, reactions
Polyamides, reactions
Polyesters, reactions
Polyurethanes, reactions
(carbide and oxycarbide based compns. and ~~nanorods~~)

IT ~~Nanotubes~~
(carbon; carbide and oxycarbide based compns. and ~~nanorods~~)
)

IT Catalyst supports
Catalysts
Hydrodesulfurization
Hydrogenolysis
(fluid phase catalytic chem. reactions; carbide and oxycarbide based compns. and ~~nanorods~~)

IT Carbides
Carbides
Oxides (inorganic)
Oxides (inorganic), uses
(oxycarbides; carbide and oxycarbide based compns. and ~~nanorods~~)

IT 7439-88-5, Iridium, uses 7439-98-7, Molybdenum, uses 7440-03-1, Niobium, uses 7440-04-2, Osmium, uses 7440-05-3, Palladium, uses 7440-06-4, Platinum, uses 7440-16-6, Rhodium, uses 7440-18-8, Ruthenium, uses 7440-25-7, Tantalum, uses 7440-32-6, Titanium, uses 7440-33-7, Tungsten, uses 7440-58-6, Hafnium, uses 7440-62-2, Vanadium, uses 7440-67-7, Zirconium, uses 12070-10-9, Vanadium carbide 12070-12-1, Tungsten carbide 12627-57-5, Molybdenum carbide 15855-70-6, Ammonium tungstate
(carbide and oxycarbide based compns. and ~~nanorods~~)

IT 409-21-2, Silicon carbide, reactions 1343-93-7, Phosphotungstic acid 9002-88-4, Polyethylene 9003-53-6, Polystyrene 9004-34-6, Cellulose, reactions 9016-00-6,

Poly(dimethylsiloxane) 12027-67-7, Ammonium molybdate 14284-90-3,
Molybdenum acetyl acetate
(carbide and oxycarbide based compns. and nanorods)
IT 124-38-9, Carbon dioxide, uses 7732-18-5, Water, uses 7782-44-7,
Oxygen, uses 10024-97-2, Nitrous oxide, uses
10102-43-9, Nitric oxide, uses 10102-44-0, Nitrogen
dioxide, uses
(oxidant; carbide and oxycarbide based compns. and nanorods
)
OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)
RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L106 ANSWER 11 OF 12 HCA COPYRIGHT 2010 ACS on STN

AN 130:29221 HCA Full-text
TI Preparation of solid porous matrixes for pharmaceutical uses
IN Unger, Evan C.
PA ImaRx Pharmaceutical Corp., USA
SO PCT Int. Appl., 139 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 6

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 9851282	A1	19981119	WO 1998-US9570	19980512
	US 20020039594	A1	20020404	US 1998-75477	19980511
	AU 9873787	A	19981208	AU 1998-73787	19980512
	EP 983060	A1	20000308	EP 1998-921109	19980512
	US 20010018072	A1	20010830	US 2001-828762	20010409
	US 20040091541	A1	20040513	US 2003-622027	20030716
PRAI	US 1997-46379P	P	19970513		
	US 1998-75477	A	19980511		
	WO 1998-US9570	W	19980512		
	US 2001-828762	B1	20010409		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A solid porous matrix formed from a surfactant, a solvent, and a bioactive
agent is described. Thus, amphotericin **nanoparticles** were prepd. by using
ZrO2 beads and a surfactant. The mixt. was milled for 24 h.
IT 9004-34-6, Cellulose, biological studies 10024-97-2,
Nitrogen oxide (N2O), biological studies

```

      (prepn. of solid porous matrixes for pharmaceutical uses)

```

RN 9004-34-6 HCA
CN Cellulose (CA INDEX NAME)
*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
RN 10024-97-2 HCA
CN Nitrogen oxide (N2O) (CA INDEX NAME)

O==N==N

IC ICM A61K009-10
 CC 63-6 (Pharmaceuticals)
 IT Drug delivery systems
 (nanoparticles; prepn. of solid porous matrixes for
 pharmaceutical uses)
 IT 9015-82-1 9028-31-3, Aldose reductase 125978-95-2, Nitric
 oxide synthase
 (inhibitors; prepn. of solid porous matrixes for
 pharmaceutical uses)
 IT 661-97-2 677-56-5, Propane-1,1,1,2,2,3-hexafluoro 678-26-2,
 Perfluoropentane 684-16-2, Hexafluoroacetone 685-63-2,
 Hexafluoro-1,3-butadiene 689-97-4, Vinyl acetylene 692-50-2,
 Hexafluoro-2-butyne 752-61-4, Digitalin 768-94-5, Amantadine
 818-92-8, 3-FluoroPropylene 846-50-4, Temazepam 921-13-1,
 Chlorodinitromethane 927-84-4, Trifluoromethyl peroxide 928-45-0,
 Butyl nitrate 968-93-4, Testolactone 987-24-6, Betamethasone
 acetate 990-73-8, Fentanyl citrate 1070-11-7, Ethambutol
 hydrochloride 1119-94-4, Lauryltrimethylammonium bromide
 1119-97-7, Myristyltrimethylammonium bromide 1172-18-5 1177-87-3,
 Dexamethasone acetate 1191-96-4, EthylCyclopropane 1306-06-5,
 Hydroxylapatite 1397-89-3, Amphotericin B 1400-61-9, Nystatin
 1404-04-2, Neomycin 1405-37-4, Capreomycin sulfate 1493-03-4,
 Difluoriodomethane 1597-82-6, Paramethasone acetate 1630-94-0,
 1,1-DimethylCyclopropane 1691-13-0, 1,2-Difluoroethylene
 1722-62-9, Mepivacaine hydrochloride 1759-88-2 1867-66-9, Ketamine
 hydrochloride 2022-85-7, Flucytosine 2068-78-2, Vincristine
 sulfate 2314-97-8, IodotriFluoromethane 2366-52-1, 1-Fluorobutane
 2375-03-3, Methylprednisolone sodium succinate 2392-39-4,
 Dexamethasone sodium phosphate 2511-95-7, 1,2-DimethylCyclopropane
 2551-62-4, Sulfur hexafluoride 3116-76-5, Dicloxacillin 3385-03-3,
 Flunisolide 3458-28-4, Mannose 3485-14-1, Cyclacillin 3511-16-8,
 Hetacillin 3529-04-2, Benzyldimethylhexadecylammonium bromide
 3810-74-0, Streptomycin sulfate 3858-89-7, Chlorprocaine
 hydrochloride 4185-80-2, Methotrimeprazine hydrochloride
 4428-95-9, Foscarnet 4431-00-9, Aurintricarboxylic acid 4697-36-3,
 Carbenicillin 4786-20-3, Crotononitrile 4901-75-1,
 3-Ethyl-3-methyldiaziridine 5534-09-8, Beclomethasone dipropionate
 5536-17-4, Arabinosyl adenine 5611-51-8, Triamcinolone hexacetonide
 5714-22-7, Sulfur fluoride (S2F10) 6000-74-4, Hydrocortisone sodium
 phosphate 7281-04-1, Benzyldimethyldodecylammonium bromide
 7297-25-8, Erythritol tetranitrate 7439-89-6, Iron, biological
 studies 7440-01-9, Neon, biological studies 7440-06-4D, Platinum,
 compds., biological studies 7440-15-5, Rhenium, biological studies
 7440-24-6, Strontium, biological studies 7440-26-8, Technetium,
 biological studies 7440-48-4, Cobalt, biological studies
 7440-63-3, Xenon, biological studies 7440-65-5, Yttrium, biological
 studies 7601-55-0, Metocurine iodide 7637-07-2, biological studies
 7647-14-5, Sodium chloride, biological studies 7681-14-3,
 Prednisolone tebutate 7727-37-9, Nitrogen, biological studies
 7728-73-6 7782-41-4, Fluorine, biological studies 7782-44-7,
 Oxygen, biological studies 7783-82-6, Tungsten hexafluoride

9001-75-6, Pepsin 9001-78-9, Alkaline phosphatase 9002-01-1, Streptokinase 9002-04-4, Thrombin 9002-60-2, Adrenocorticotrophic hormone, biological studies 9002-61-3 9002-72-6, Growth hormone 9002-79-3, Melanocyte stimulating hormone 9002-89-5, Poly(vinyl alcohol) 9003-11-6 9003-39-8, PVP 9004-10-8, Insulin, biological studies ~~9004-34-6~~, Cellulose, biological studies 9004-54-0, Dextran, biological studies 9004-61-9, Hyaluronic acid 9004-67-5, Methyl Cellulose 9005-25-8, Starch, biological studies 9005-27-0, HETA-starch 9005-32-7, Alginic acid 9005-49-6, Heparin, biological studies 9005-64-5, Polyoxyethylene sorbitan monolaurate 9005-65-6, Polyoxyethylene sorbitan monooleate 9005-66-7, Polyoxyethylene sorbitan monopalmitate 9005-67-8, Polyoxyethylene sorbitan monostearate 9005-71-4, Polyoxyethylene sorbitan tristearate 9007-12-9, Calcitonin 9007-92-5, Glucagon, biological studies 9011-14-7, PMMA 9011-97-6, Cholecystokinin 9015-68-3, Asparaginase 9015-71-8, Corticotropin releasing factor 9036-19-5, Octoxynol 9039-53-6, Urokinase 9061-61-4, Nerve growth factor ~~10024-97-2~~, Nitrogen oxide (N₂O), biological studies 11000-17-2, Vasopressin 11056-06-7, Bleomycin 11096-26-7, Erythropoietin 13264-41-0, Cetyldimethylethylammonium chloride 13292-46-1, Rifampin 13311-84-7, Flutamide 13647-35-3, Trilostane 15500-66-0, Pancuronium bromide 15663-27-1, Cisplatin 15686-71-2, Cephalixin 15687-27-1, Ibuprofen 16009-13-5, Hemin 16136-85-9 17598-65-1, Deslanoside 18010-40-7, Bupivacaine hydrochloride 18323-44-9, Clindamycin 18378-89-7, Plicamycin 18773-88-1, Benzyldimethyltetradecylammonium bromide 20187-55-7, Bendazac 20274-91-3 20830-75-5, Digoxin 21829-25-4, Nifedipine 22204-53-1, Naproxen 22494-42-4, Diflunisal 22916-47-8, Miconazole 23110-15-8, Fumagillin 23541-50-6, Daunorubicin hydrochloride 24356-66-9 24764-97-4, 2-Bromobutyraldehyde 24991-23-9 25104-18-1, Polylysine 25151-81-9, Prostanoid acid 25316-40-9, Adriamycin 25322-68-3 25322-68-3D, PEG, ethers 25322-69-4, Polypropylene glycol 25513-46-6, Polyglutamic acid 26023-30-3, Poly[oxy(1-methyl-2-oxo-1,2-ethanediyl)] 26100-51-6, Poly(lactic acid) 26171-23-3, Tolmetin 26780-50-7, Glycolide-lactide copolymer 26787-78-0, Amoxicillin 26839-75-8, Timolol 28911-01-5, Triazolam 29121-60-6, Vaninolol 29767-20-2, Teniposide 30516-87-1, Azidothymidine 31637-97-5, Etofibrate 33069-62-4, Taxol 33125-97-2, Etomidate 33419-42-0, Etoposide 33507-63-0, Substance p 34077-87-7, DiChlorotrifluoroethane 34787-01-4, Ticarcillin 36322-90-4 36637-19-1, Etidocaine hydrochloride 36791-04-5, Ribavirin 38000-06-5, Polylysine 38194-50-2, Sulindac 38821-53-3, Cephadrine 39391-18-9, Cyclooxygenase 41575-94-4, Carboplatin 42399-41-7, Diltiazem 47141-42-4, Levobunolol 50370-12-2, Cefadroxil 50402-72-7, Piperidine-2,3,6-trimethyl 50700-72-6, Vecuronium bromide 50972-17-3, Bacampicillin 51264-14-3, Amsacrine 52205-73-9, Estramustine phosphate sodium 52365-63-6, Dipivefrin 53045-71-9, 1-Pentene-3-bromo 53188-07-1, Trolox 53678-77-6, Muramyl dipeptide 53994-73-3, Cefaclor 54965-24-1, Tamoxifen citrate 55142-85-3, Ticlopidine 57223-18-4, 1-Nonen-3-yne 59277-89-3, Acyclovir 59467-96-8, Midazolam

hydrochloride 60118-07-2, Endorphin 62031-54-3, Fibroblast growth factor 62229-50-9, Epidermal growth factor 62232-46-6, Bifemelane hydrochloride 62571-86-2, Captopril 62683-29-8, Colony stimulating factor 63659-18-7, Betaxolol 65277-42-1, Ketoconazole 68302-57-8 68367-52-2, Sorbinil 69279-90-9, Ansamitocin 72702-95-5, Ponalrestat 73218-79-8, Apraclonidine hydrochloride 73984-11-9 74381-53-6, Leuprolide acetate 74790-08-2, Spiroplatin 75847-73-3, Enalapril 76547-98-3, Lisinopril 77181-69-2, Sorivudine 80755-87-9 81486-22-8, Nipradilol 82159-09-9, Epalrestat 82410-32-0, Ganciclovir 82964-04-3, Tolrestat 83869-56-1, Granulocyte macrophage colony stimulating factor 86090-08-6, Angiostatin 88096-12-2 89149-10-0, 15-Deoxyspergualin 98023-09-7 99896-85-2 106956-32-5, Oncostatin M 113852-37-2, Cidofovir 116632-15-6, 1.2.3-Nonadecanetricarboxylic acid 2-hydroxytrimethylester 119813-10-4, Carzelesin 120279-96-1, Dorzolamide 120287-85-6D, Cetrorelix, derivs. 121181-53-1, Filgrastim 124389-07-7, Muramyltripeptide 127464-60-2, Vascular endothelial growth factor 127984-74-1, Somatuline 130209-82-4, Latanoprost 139639-23-9, Tissue plasminogen activator 141436-78-4, Protein kinase c

(~~prepn.~~ of solid porous matrixes for pharmaceutical uses)

OSC.G 17 THERE ARE 17 CAPLUS RECORDS THAT CITE THIS RECORD (17 CITINGS)

RE.CNT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L106 ANSWER 12 OF 12 HCA COPYRIGHT 2010 ACS on STN

AN 116:231308 HCA Full-text

OREF 116:39063a,39066a

TI Photolytic interface for HPLC-chemiluminescence detection of nonvolatile N-nitroso compounds

IN Conboy, James J.; Hotchkiss, Joseph H.

PA Cornell Research Foundation, Inc., USA

SO U.S., 12 pp.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 5094815	A	19920310	US 1988-195923	19880518
	US 5366900	A	19941122	US 1993-10578	19930128
PRAI	US 1988-195923	A3	19880518		
	US 1991-798490	B1	19911224		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Described are a photolytic interface app. and its use in series between a HPLC and a chemiluminescence detector for the detection of trace (~~nanogram~~) amts. of N-nitroso compds. including N-nitrosamides and non-volatile N-nitrosamines in aq.-based fluid samples. HPLC effluent contg. sepd. N-nitrosoamino acids and N-nitrosoamino amides is introduced into a glass coil with a purge stream of He and irradiated with UV light. NO cleaved by photolysis is rapidly sepd. from solvent through a series of cold traps and


carried by the He into the reaction chamber of a chemiluminescence detector. Biol. matrixes, such as urine and gastric fluid, can be analyzed directly at high sensitivity without concn. and/or extn. Figures show diagrams of the photolytic interface app. and the total app. system incorporating the interface app. as well as chromatograms showing resoln. of std.N-nitroso compds. in std. solns. and in urine and porcine gastric juice samples.

IT 70-25-7

(molar response ratios of, in analyzer having HPLC and photolytic interface and chemiluminescence detector)

RN 70-25-7 HCA

CN Guanidine, N-methyl-N'-nitro-N-nitroso- (CA INDEX NAME)



IC ICM G01N021-76

INCL 422052000

CC 9-1 (Biochemical Methods)

Section cross-reference(s): 80

IT Helium-group gases, uses

(in photolytic interface app. for carrying **nitric oxide formed** from HPLC-sepd. N-nitroso compds. to chemiluminescence detector)

IT 7440-59-7, Helium, uses

(in photolytic interface app. for carrying **nitric oxide formed** from HPLC-sepd. N-nitroso compds. to chemiluminescence detector)

IT 62-75-9, NDMA 70-25-7 684-93-5 759-73-9 3475-63-6,
N-Nitrosotrimethylurea 7519-36-0, N-Nitrosoproline 13256-22-9,
N-Nitrososarcosine 30310-80-6, N-Nitrosohydroxyproline 88381-44-6
103659-08-1

(molar response ratios of, in analyzer having HPLC and photolytic interface and chemiluminescence detector)

OSC.G 23 THERE ARE 23 CAPLUS RECORDS THAT CITE THIS RECORD (24 CITINGS)

RE.CNT 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> D L107 1-3 BIB ABS HITSTR HITIND

L107 ANSWER 1 OF 3 HCA COPYRIGHT 2010 ACS on STN

AN 142:435873 HCA Full-text

TI A medical device with ~~nanofiber~~ outer surface layer incorporating nitric oxide and poly(ethylenimine) ~~diazoniumdiolate~~ for insertion in to vascular system

IN Andersen, Erik; Smith, Daniel; Reneker, Darrell

PA Cube Medical A/S, Den.; The University of Akron

SO PCT Int. Appl., 21 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 3

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2005039664	A2	20050506	WO 2004-US33949	20041014
	WO 2005039664	A3	20050630		
	EP 1691856	A2	20060823	EP 2004-795149	20041014
	CN 1874799	A	20061206	CN 2004-80032370	20041014
	JP 2008539807	T	20081120	JP 2006-535667	20041014
	US 20070207179	A1	20070906	US 2006-595339	20061229
PRAI	DK 2003-1514	A	20031014		
	US 2003-510520P	P	20031014		
	DK 2003-1864	A	20031216		
	US 2003-529629P	P	20031216		
	DK 2004-671	A	20040429		
	US 2004-566087P	P	20040429		
	WO 2004-US33949	W	20041014		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A medical device, such as a guide wire, an embolization device, or a guide shaft for a micro-catheter, comprises a solid and/or non-expandable core member made from e.g. metal, such as tantalum, and an outer surface layer, which is formed by electrospun **nanofibers**. The outer surface layer may incorporate a pharmaceutically active substance, such as a nitric oxide (NO) donor for release in the vascular or neurovascular system of a living being. The NO donor may be incorporated in a polymer, such as a polymeric linear poly(ethylenimine) **diazoniumdiolate**.

IT 10102-43-9, Nitric oxide, biological studies
(medical device with **nanofiber** outer surface layer
incorporating nitric oxide and poly(ethylenimine)
diazoniumdiolate for insertion in to vascular system)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

IC ICM A61L029-00

CC 63-7 (Pharmaceuticals)

ST nitric oxide donor polyethylenimine **nanofiber** coating
medical device; polyethylenimine **diazoniumdiolate** acid
nanofiber medical embolization device

IT Embolism
(embolization; medical device with **nanofiber** outer
surface layer incorporating nitric oxide and poly(ethylenimine)
diazoniumdiolate for insertion in to vascular system)

IT Polyesters, biological studies
(lactide; medical device with **nanofiber** outer surface

layer incorporating nitric oxide and poly(ethylenimine)
diazeniumdiolate for insertion in to vascular system)

IT Coating materials
Drug delivery systems
Drugs
Nanofibers
(medical device with **nanofiber** outer surface layer
incorporating nitric oxide and poly(ethylenimine)
diazeniumdiolate for insertion in to vascular system)

IT Acids, biological studies
Collagens, biological studies
Fluoropolymers, biological studies
Polyurethanes, biological studies
Synthetic fibers
(medical device with **nanofiber** outer surface layer
incorporating nitric oxide and poly(ethylenimine)
diazeniumdiolate for insertion in to vascular system)

IT Synthetic polymeric fibers, biological studies
(polyethylenimine; medical device with **nanofiber** outer
surface layer incorporating nitric oxide and poly(ethylenimine)
diazeniumdiolate for insertion in to vascular system)

IT Medical goods
(stents; medical device with **nanofiber** outer surface
layer incorporating nitric oxide and poly(ethylenimine)
diazeniumdiolate for insertion in to vascular system)

IT Medical goods
(wires; medical device with **nanofiber** outer surface layer
incorporating nitric oxide and poly(ethylenimine)
diazeniumdiolate for insertion in to vascular system)

IT 10102-43-9, Nitric oxide, biological studies
(medical device with **nanofiber** outer surface layer
incorporating nitric oxide and poly(ethylenimine)
diazeniumdiolate for insertion in to vascular system)

IT 7440-25-7, Tantalum, biological studies 9002-84-0,
Polytetrafluoroethylene 26023-30-3,
Poly[oxy(1-methyl-2-oxo-1,2-ethanediyl)] 26680-10-4, Polylactide
(medical device with **nanofiber** outer surface layer
incorporating nitric oxide and poly(ethylenimine)
diazeniumdiolate for insertion in to vascular system)

IT 9002-98-6D, diazenium diolate derivs. 26913-06-4D,
Poly[imino(1,2-ethanediyl)], diazenium diolate derivs.
(**nanofiber**; medical device with **nanofiber** outer
surface layer incorporating nitric oxide and poly(ethylenimine)
diazeniumdiolate for insertion in to vascular system)

OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L107 ANSWER 2 OF 3 HCA COPYRIGHT 2010 ACS on STN

AN 138:381586 HCA Full-text

TI Superoxide-dependent consumption of nitric oxide in biological media
may confound in vitro experiments

AU Keynes, Robert G.; Griffiths, Charmaine; Garthwaite, John
 CS Cruciform Building, Wolfson Institute for Biomedical Research,
 University College London, London, WC1E 6BT, UK
 SO Biochemical Journal (2003), 369(2), 399-406
 CODEN: BIJOAK; ISSN: 0264-6021
 PB Portland Press Ltd.
 DT Journal
 LA English
 AB NO functions ubiquitously as a biol. messenger but was also implicated in various pathologies, a role supported by many reports that exogenous or endogenous NO can kill cells in tissue culture. In the course of expts. aimed at examng. the toxicity of exogenous NO towards cultured cells, the authors found that most of the NO delivered using a NONOate (diazoniumdiolate) donor was removed by reaction with the tissue-culture medium. Two NO-consuming ingredients were identified: Hepes buffer and, under lab. lighting, the vitamin riboflavin. In each case, the loss of NO was reversed by the addn. of superoxide dismutase. The effect of Hepes was obsd. over a range of NONOate concns. (producing up to 1 μ M NO). Furthermore, from measurements of sol. guanylate cyclase activity, Hepes-dependent NO consumption remained significant at the low nanomolar NO concns. relevant to physiol. NO signaling. The combination of Hepes and riboflavin (in the light) acted synergistically to the extent that, instead of a steady-state concn. of about 1 μ M being generated, NO was undetectable (<10 nM). Again, the consumption could be inhibited by superoxide dismutase. A scheme is proposed whereby a 'vicious cycle' of superoxide radical ($O\bullet-2$) formation occurs as a result of oxidn. of Hepes to its radical species, fuelled by the subsequent reaction of $O\bullet-2$ with NO to form peroxynitrite ($ONOO-$). The inadvertent prodn. of $ONOO-$ and other reactive species in biol. media, or the assocd. loss of NO, may contribute to the adverse effects, or otherwise, of NO in vitro.
 IT 10102-43-9, Nitric oxide, biological studies
 (superoxide, riboflavin, and buffer effect on NO consumption in biol. media)
 RN 10102-43-9 HCA
 CN Nitrogen oxide (NO) (CA INDEX NAME)

$N=O$

CC 9-11 (Biochemical Methods)
 IT 77-86-1, Tris buffer 83-88-5, Riboflavin, biological studies
 7365-45-9, HEPES 9054-89-1, Superoxide dismutase 10102-43-9
 , Nitric oxide, biological studies 11062-77-4, Superoxide
 (superoxide, riboflavin, and buffer effect on NO consumption in biol. media)
 OSC.G 27 THERE ARE 27 CAPLUS RECORDS THAT CITE THIS RECORD (27 CITINGS)
 RE.CNT 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L107 ANSWER 3 OF 3 HCA COPYRIGHT 2010 ACS on STN

AN 134:300856 HCA Full-text

TI Nitric oxide-modified linear poly(ethylenimine) fibers for coating of medical devices

IN Smith, Daniel; Reneker, Darrell

PA University of Akron, USA

SO PCT Int. Appl., 13 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2001026702	A2	20010419	WO 2000-US27769	20001006
	WO 2001026702	A3	20011213		
	US 6737447	B1	20040518	US 2000-571444	20000516
	CA 2386765	A1	20010419	CA 2000-2386765	20001006
	EP 1220694	A2	20020710	EP 2000-970658	20001006
	EP 1220694	B1	20030416		
	AT 237372	T	20030515	AT 2000-970658	20001006
	US 20040131753	A1	20040708	US 2003-738582	20031216
	US 6855366	B2	20050215		
PRAI	US 1999-158673P	P	19991008		
	US 2000-571444	A	20000516		
	WO 2000-US27769	W	20001006		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A novel coating for medical devices provides nitric oxide delivery using **nanofibers** of linear poly(ethylenimine) **diazeniumdiolate**. Linear poly(ethylenimine) **diazeniumdiolate** releases nitric oxide (NO) in a controlled manner to tissues and organs to aid the healing process and to prevent injury to tissues at risk of injury. Electrospun **nanofibers** of linear poly(ethylenimine)**diazeniumdiolate** deliver therapeutic levels of NO to the tissues surrounding a medical device while minimizing the alteration of the properties of the device. A **nanofiber** coating, because of the small size and large surface area per unit mass of the **nanofibers**, provides a much larger surface area per unit mass while minimizing changes in other properties of the device (no data).

IT ~~10102-43-9~~, Nitric oxide, biological studies
(nitric oxide-modified linear poly(ethylenimine) fibers for coating of medical devices)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

N==O

IC ICM A61L027-00

CC 63-7 (Pharmaceuticals)

Section cross-reference(s): 38

IT Synthetic fibers
(~~nano~~-; nitric oxide-modified linear poly(ethylenimine)
fibers for coating of medical devices)
IT 10102-43-9, Nitric oxide, biological studies
(nitric oxide-modified linear poly(ethylenimine) fibers for coating
of medical devices)
OSC.G 16 THERE ARE 16 CAPLUS RECORDS THAT CITE THIS RECORD (16
CITINGS)
RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> D L111 1 BIB ABS HITSTR HITIND

L111 ANSWER 1 OF 1 HCA COPYRIGHT 2010 ACS on STN
AN 134:67585 HCA Full-text
TI Tyrosine nitration by peroxynitrite ~~formed~~ from
~~nitric oxide~~ and superoxide generated by
xanthine oxidase
AU Sawa, Tomohiro; Akaike, Takaaki; Maeda, Hiroshi
CS Department of Microbiology, Kumamoto University School of Medicine,
Kumamoto, 860-0811, Japan
SO Journal of Biological Chemistry (2000), 275(42), 32467-32474
CODEN: JBCHA3; ISSN: 0021-9258
PB American Society for Biochemistry and Molecular Biology
DT Journal
LA English
AB Peroxynitrite (ONOO-) is a potent nitrating and oxidizing agent that is
formed by a rapid reaction of nitric oxide (NO) with superoxide anion (O2).
It appears to be involved in the pathophysiol. of many inflammatory and
neurodegenerative diseases. It has recently been reported that ONOO-
generated at neutral pH from NO and O2 (NO/O2) was substantially less
efficient than preformed ONOO- at nitrating tyrosine. Here we re-evaluated
tyrosine nitration by NO/O2 with a shorter incubation period and a more
sensitive electrochem. detection system. Appreciable amts. of nitrotyrosine
were produced by ONOO- formed in situ (2.9 μ M for 5 min; 10 nM/s) by NO/O2
flux obtained from propylamine ~~NONOate~~ (CH3N[N(O)NO]- (CH2)3NH2+CH3) and
xanthine oxidase using pterin as a substrate in phosphate buffer (pH 7.0)
contg. 0.1 mM L-tyrosine. The yield of nitrotyrosine by this NO/O2 flux was
approx. 70% of that produced by the same flux of preformed ONOO- (2.9 μ M/5
min). When hypoxanthine was used as a substrate, tyrosine nitration by
NO/O2 was largely eliminated because of the inhibitory effect of uric acid
produced during the oxidn. of hypoxanthine. Tyrosine nitration caused by
NO/O2 was inhibited by the ONOO- scavenger ebselen and was enhanced 2-fold
by NaHCO3, as would be expected, because CO2 promotes tyrosine nitration.
The profile of nitrotyrosine and dityrosine formation produced by NO/O2 flux
(2.9 μ M/5 min) was consistent with that produced by preformed ONOO-.
Tyrosine nitration predominated compared with dityrosine formation caused by
a low ~~nanomolar~~ flux of ONOO- at physiol. concns. of free tyrosine (<0.5
mM). In conclusion, our results show that NO generated with O2 nitrates
tyrosine with a reactivity and efficacy similar to those of chem.

synthesized ONOO-, indicating that ONOO- can be a significant source of tyrosine nitration in physiol. and pathol. events in vivo.

IT 10102-43-9, Nitric oxide, biological studies
19059-14-4, Peroxynitrite
(tyrosine nitration by peroxynitrite formed from
nitric oxide and superoxide generated
by xanthine oxidase)


RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



RN 19059-14-4 HCA

CN Peroxynitrite (8CI, 9CI) (CA INDEX NAME)



CC 6-1 (General Biochemistry)

IT Nitration
(tyrosine nitration by peroxynitrite formed from
nitric oxide and superoxide generated
by xanthine oxidase)

IT 9002-17-9, Xanthine oxidase 10102-43-9, Nitric oxide,
biological studies 11062-77-4, Superoxide 19059-14-4,
Peroxynitrite
(tyrosine nitration by peroxynitrite formed from
nitric oxide and superoxide generated
by xanthine oxidase)

IT 60-18-4, L-Tyrosine, biological studies
(tyrosine nitration by peroxynitrite formed from
nitric oxide and superoxide generated
by xanthine oxidase)

OSC.G 94 THERE ARE 94 CAPLUS RECORDS THAT CITE THIS RECORD (94
CITINGS)

RE.CNT 51 THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

CLAIM 1 AND RELATED

=> D L75 1-20 BIB ABS HITSTR HITIND


L75 ANSWER 1 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 142:214882 HCA Full-text
 TI Stabilization and ionic triggering of nitric oxide release
 IN Smith, Daniel J.
 PA The University of Akron, USA
 SO PCT Int. Appl., 24 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2005011575	A2	20050210	WO 2004-US23867	20040726
	WO 2005011575	A3	20060112		
	EP 1648527	A2	20060426	EP 2004-779101	20040726
	US 20090136410	A1	20090528	US 2007-565573	20070226
PRAI	US 2003-490218P	P	20030725		
	WO 2004-US23867	W	20040726		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Provided is a method for producing nitric oxide that employs an ion exchange resin. Also provided is a method for producing nitric oxide that combines a salt with a gel or cream. A method is provided for producing nitric oxide that combines a pH adjuster with a diazeniumdiolate-contg. compd. or compn.
 IT 113-21-3, Lactate, analysis
 (stabilization and ionic triggering of nitric oxide release)
 RN 113-21-3 HCA
 CN Propanoic acid, 2-hydroxy-, ion(1-) (CA INDEX NAME)



IT 10102-43-9, Nitric oxide, biological studies
 (stabilization and ionic triggering of nitric oxide release)
 RN 10102-43-9 HCA
 CN Nitrogen oxide (NO) (CA INDEX NAME)



IT 201168-09-4D, Dowex 1X400, reaction with NONOates
 (stabilization and ionic triggering of nitric oxide release)
 RN 201168-09-4 HCA
 CN Dowex 1X400 (9CI) (CA INDEX NAME)
 *** STRUCTURE DIAGRAM IS NOT AVAILABLE ***
 IC ICM A61K
 CC 9-16 (Biochemical Methods)
 IT Ion exchangers

Nanofibers
Nanoparticles
pH

(stabilization and ionic triggering of nitric oxide release)

IT 113-21-3, Lactate, analysis 126-44-3, Citrate, analysis 14265-44-2, Phosphate, analysis
(stabilization and ionic triggering of nitric oxide release)
IT 10102-43-9, Nitric oxide, biological studies
(stabilization and ionic triggering of nitric oxide release)
IT 16545-40-7 27561-78-0 201168-09-40, Dowex 1X400, reaction with NONOates 839676-39-0 839676-40-3 839676-41-4
(stabilization and ionic triggering of nitric oxide release)
OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)
RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L75 ANSWER 2 OF 20 HCA COPYRIGHT 2010 ACS on STN
AN 141:370637 HCA Full-text
TI Fibrous assemblies that sequester reactive materials
IN Reneker, Darrell H.; Smith, Daniel J.
PA The University of Akron, USA
SO PCT Int. Appl., 15 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	WO 2004094050	A2	20041104	WO 2004-US12673	20040423
	WO 2004094050	A3	20050414		
	AU 2004233347	A1	20041104	AU 2004-233347	20040423
	CA 2523957	A1	20041104	CA 2004-2523957	20040423
	EP 1624953	A2	20060215	EP 2004-760164	20040423
	JP 2006525445	T	20061109	JP 2006-513289	20040423
	CN 1917836	A	20070221	CN 2004-80017111	20040423
	IN 2005DN05060	A	20071005	IN 2005-DN5060	20051107
	IN 235762	A1	20090904		
	US 20060280781	A1	20061214	US 2006-554191	20060803
	IN 2009DN03391	A	20100409	IN 2009-DN3391	20090525
PRAI	US 2003-464879P	P	20030423		
	WO 2004-US12673	W	20040423		
	IN 2005-DN5060	A3	20051107		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A fibrous assembly is provided for performing site-specific chem. In general the present invention provides a fibrous assembly comprising a first fiber that sequesters a first reactive component; and a second fiber that sequesters a second reactive component, wherein at least the first or second fiber releases its reactive component when the fiber is in the presence of a releasing agent, and wherein when the at least first or second fiber releases its reactive component, the first and second reactive components react with each other to form a reaction product. Related methods of manuf.

and use are also provided. For example, a nanofiber assembly was prepd. contg. two types of fibers, each sequestering a reactive component: fiber one contained ascorbic acid and fiber two contained potassium nitrite. When exposed to moisture, the assembly releases ingredients to give ascorbic acid and NO₂⁻, which react to form nitric oxide. Alternatively, nitrite and/or ascorbic acid may be immobilized such as by being adsorbed onto an ~~ion~~ exchange resin bead, which is then incorporated into polymer fibers or nanofibers. Fiber assemblies as described above are envisioned as being used in nitric oxide-releasing medical dressings for the treatment of wounds and other lesions of the skin such as warts. This method may also be useful in other fields where the sequestration of reactive component is desired, such as in the creation of epoxy-type adhesives.

IT 10102-43-9, Nitric oxide, formation (nonpreparative)
(fibrous assemblies that sequester reactive materials for delivery to targeted locations)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

IC ICM B01F
CC 63-7 (Pharmaceuticals)
ST fiber sequestrant reactive chem wound dressing adhesive; nitric oxide ~~ascorbate~~ nitrite nanofiber assembly
IT Spheres
(beads, ~~ion exchangers~~; fibrous assemblies that sequester reactive materials for delivery to targeted locations)
IT ~~Ion exchangers~~
(beads; fibrous assemblies that sequester reactive materials for delivery to targeted locations)
IT 10102-43-9, Nitric oxide, formation (nonpreparative)
(fibrous assemblies that sequester reactive materials for delivery to targeted locations)
RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L75 ANSWER 3 OF 20 HCA COPYRIGHT 2010 ACS on STN
AN 140:14124 HCA Full-text
TI Purification and characterization of a ubiquitin-like peptide with macrophage stimulating, antiproliferative and ribonuclease activities from the mushroom *Agrocybe cylindracea*
AU Ngai, Patrick H. K.; Wang, H. X.; Ng, T. B.
CS Faculty of Medicine, Department of Biochemistry, The Chinese University of Hong Kong, Shatin, Hong Kong
SO Peptides (New York, NY, United States) (2003), 24(5), 639-645
CODEN: PPTDD5; ISSN: 0196-9781
PB Elsevier Science Inc.
DT Journal

LA English

AB A peptide, with a mol. mass of 9.5 kDa and demonstrating an N-terminal sequence similar to ubiquitin, was isolated from fruiting bodies of the mushroom *Agrocybe cylindracea*. The peptide was isolated with a purifn. protocol involving ion exchange chromatog. on DEAE-cellulose, affinity chromatog. on Affi-gel blue gel, FPLC- ion exchange chromatog. on Mono S and FPLC-gel filtration on Superdex 75. The peptide was unadsorbed on DEAE-cellulose and adsorbed on Affi-gel blue gel and Mono S. It showed antiproliferative activity on leukemia cell line (M1) and hepatoma cell line (HepG2), and enhanced nitric oxide prodn. in murine peritoneal macrophages with a potency comparable to that of lipopolysaccharide. A pH of 6.0 was required for optimal RNase activity. Its RNase activity was stable over the temp. range of 0-60°. It exerted ribonucleolytic activity preferentially on polyC, much lower activity on polyU, and negligible activity on polyA and polyG.

IT 10102-43-9, Nitric oxide, biological studies
 (prodn. of, effect of isolated ubiquitin-like peptide on; ubiquitin-like peptide from the mushroom *Agrocybe cylindracea* with macrophage stimulating, antiproliferative, and RNase activities)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

$\text{N}=\text{O}$

CC 6-3 (General Biochemistry)
 Section cross-reference(s): 10

IT 10102-43-9, Nitric oxide, biological studies
 (prodn. of, effect of isolated ubiquitin-like peptide on; ubiquitin-like peptide from the mushroom *Agrocybe cylindracea* with macrophage stimulating, antiproliferative, and RNase activities)

OSC.G 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD (8 CITINGS)

RE.CNT 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L75 ANSWER 4 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 138:226775 HCA Full-text

TI Preparation of morpholinosydnonimine-sugar conjugates as nitric oxide donors

IN Wang, Peng George; Wu, Xuejun; Tang, Xiaoping

PA Wayne State University, USA

SO U.S. Pat. Appl. Publ., 10 pp.
 CODEN: USXXCO

DT Patent

LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----

PI US 20030050256 A1 20030313 US 2001-925816 20010809
US 6867194 B2 20050315
PRAI US 2001-925816 20010809

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OS MARPAT 138:226775

AB Sugar-modified SIN-1 compns. are provided. The compns. are useful for generating NO in response to hydrolytic activity of a glycosidase specific for the O-glycosidic bond between the sugar and SIN-1 moieties. Pharmaceutical compns. contg. the sugar-modified SIN-1 compns. and methods of using the compns. are also provided. 3-Morpholinosydnonimine-HCl was prepd. by a std. method. To a soln. of 4-nitrophenyl (2,3,4,6-tetra-O-acetyl- α/β -D-glucopyranosyl) carbonate in anhyd. pyridine was added the above compd. The solvent was removed in vacuo to give a sticky oil and the residue was purified by silica gel column chromatog. to give a mixt. of α - and β -anomers of the morpholinosydnonimine-glucose conjugate. The mixt. was treated with NaOCH₃ in anhyd. MeOH and Amberlyst-15 ion-exchange resin was added to neutralize the reaction mixt.

IT 10102-43-9, Nitric oxide, biological studies
(prepn. of morpholinosydnonimine-sugar conjugates as nitric oxide donors)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



IC ICM A61K031-706

ICS C07H017-02; C07H019-048

INCL 514043000; 514023000; 536028100; 536017400

CC 63-6 (Pharmaceuticals)

Section cross-reference(s): 1, 33

IT Named reagents and solutions

(Ringer's lactate, liq. carrier; prepn. of morpholinosydnonimine-sugar conjugates as nitric oxide donors)

IT 9032-92-2, Glycosidase 10102-43-9, Nitric oxide, biological studies 11062-77-4, Superoxide 19059-14-4, Peroxynitrite
(prepn. of morpholinosydnonimine-sugar conjugates as nitric oxide donors)

RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L75 ANSWER 5 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 138:2681 HCA Full-text

TI L-arginine potentiates negative inotropic and metabolic effects to myocardium partly through the amiloride sensitive mechanism

AU Takeuchi, Koh; Simplaceanu, Elena; McGowan, Francis X., Jr.; Tsushima, Takao; del Nido, Pedro J.

CS Department of Cardiac Surgery, Children's Hospital, Boston and Harvard Medical School, Boston, MA, 02115, USA

SO Japanese Journal of Physiology (2002), 52(2), 207-215
 CODEN: JJPHAM; ISSN: 0021-521X

PB Center for Academic Publications Japan

DT Journal

LA English

AB Recently, cytokines have been proposed to cause cellular injury by nitric oxide (NO \cdot) mediated pathway and L-arginine has been proposed to impair intracellular pH (pHi) regulation via vacuolar type H $^{+}$ -ATPase in macrophage. We conducted this investigation on Langendorff perfused hearts of rabbits to elucidate the mechanisms involving the NO \cdot precursor L-arginine on myocardial contractile function, glycolysis, mitochondrial respiration, and intracellular alkalinization and tested the effects of amiloride. L-Arginine caused a significant loss of contractile function (96 \pm 4 mmHg in control, 53 \pm 16 during L-arginine perfusion, p<0.01) and a significant increase of pH $_i$ (7.01 \pm 0.02 prearginine infusion, 7.08 \pm 0.03 at the end of L-arginine infusion, p<0.01) along with decreased oxygen consumption (MVO $_2$) (0.94 \pm 0.32 mL/min/g dry wt.), increased lactate release, and a loss of creatine phosphate (15% loss). Amiloride could prevent the cell alkalinization and contractile dysfunction, but not the derangement of oxidative metab. caused by L-arginine in myocytes. We conclude that L-arginine has two distinct effects upon the myocardium: (1) an amiloride-sensitive loss of contractile function assocd. with intracellular alkalinization; and (2) an amiloride-insensitive inhibition of oxidative metab., possibly because of increased myocardial NO prodn.

IT 10102-43-9, Nitric oxide, biological studies
 (effect of NO precursor L-arginine on amiloride-sensitive Na $^{+}$ /H $^{+}$ exchange, myocardial contractility, oxidative metab., high-energy phosphates, glycolysis and intracellular pH)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

CC 13-6 (Mammalian Biochemistry)

IT Transport proteins
 (hydrogen ion-sodium exchanger; effect of NO precursor L-arginine on amiloride-sensitive Na $^{+}$ /H $^{+}$ exchange, myocardial contractility, oxidative metab., high-energy phosphates, glycolysis and intracellular pH)

IT 50-99-7, D-Glucose, biological studies 67-07-2, Creatine phosphate
 74-79-3, L-Arginine, biological studies 10102-43-9, Nitric oxide, biological studies 12408-02-5, Hydrogen ion, biological studies 14265-44-2, Phosphate, biological studies
 (effect of NO precursor L-arginine on amiloride-sensitive Na $^{+}$ /H $^{+}$ exchange, myocardial contractility, oxidative metab., high-energy phosphates, glycolysis and intracellular pH)

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

RE.CNT 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L75 ANSWER 6 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 135:195359 HCA Full-text

TI Preparation of cycloalkanone from cycloalkyl ~~nitrite~~

IN Yamamoto, Shoji; Sugimoto, Tsunemi

PA Ube Industries, Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	JP 2001240574	A	20010904	JP 2000-53175	20000229
PRAI	JP 2000-53175		20000229		

OS CASREACT 135:195359; MARPAT 135:195359

AB Cycloalkanone is prepd. by contact reaction of cycloalkyl ~~nitrite~~ in the presence of solid acid catalyst (with recovering the resulting NO for recycling). Thus, cyclohexyl ~~nitrite~~ was treated with NH₄-ZSM-5 in MeCN at 85° for 2 h to give 50:50 cyclohexanone and cyclohexanol with 55% conversion.

IT ~~10102-43-9P~~, Nitrogen monoxide, preparation
(prepn. of cycloalkanone from cycloalkyl ~~nitrite~~ with solid acid catalysts)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

IC ICM C07C049-403

ICS B01J021-12; B01J021-16; B01J029-10; B01J029-40; C07B061-00;
C07C045-32

CC 24-5 (Alicyclic Compounds)

ST cycloalkanone prepn solid acid catalyst; cyclohexanone prepn zeolite catalyst; cyclohexyl ~~nitrite~~ disproportionation zeolite catalyst

IT Zeolite ZSM-5
(ammonium-substituted; prepn. of cycloalkanone from cycloalkyl ~~nitrite~~ with solid acid catalysts)

IT Ketones, preparation
(cycloalkanones; prepn. of cycloalkanone from cycloalkyl ~~nitrite~~ with solid acid catalysts)

IT Zeolite NaY
(iron-substituted; prepn. of cycloalkanone from cycloalkyl ~~nitrite~~ with solid acid catalysts)

IT Cycloalkanols
(~~nitrites~~; prepn. of cycloalkanone from cycloalkyl ~~nitrite~~ with solid acid catalysts)

IT Acids, uses

(oxo; prepn. of cycloalkanone from cycloalkyl nitrite with solid acid catalysts)

IT Disproportionation catalysts

Ion exchangers

(prepn. of cycloalkanone from cycloalkyl nitrite with solid acid catalysts)

IT Acids, uses

Clay minerals

Zeolite HZSM-5

Zeolites (synthetic), uses

(prepn. of cycloalkanone from cycloalkyl nitrite with solid acid catalysts)

IT 1724-39-6P, Cyclododecanol

(prepn. of cycloalkanone from cycloalkyl nitrite with solid acid catalysts)

IT 108-93-0P, Cyclohexanol, preparation 10102-43-9P, Nitrogen monoxide, preparation

(prepn. of cycloalkanone from cycloalkyl nitrite with solid acid catalysts)

IT 1318-93-0, Montmorillonite, uses 7631-86-9, Silica, uses

(prepn. of cycloalkanone from cycloalkyl nitrite with solid acid catalysts)

IT 108-94-1P, Cyclohexanone, preparation 830-13-7P, Cyclododecanone

(prepn. of cycloalkanone from cycloalkyl nitrite with solid acid catalysts)

IT 50744-58-6, Cyclododecyl nitrite

(prepn. of cycloalkanone from cycloalkyl nitrite with solid acid catalysts)

IT 5156-40-1P, Cyclohexyl nitrite

(prepn. of cycloalkanone from cycloalkyl nitrite with solid acid catalysts)

L75 ANSWER 7 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 134:160771 HCA Full-text

TI Nitrite uptake and metabolism and oxidant stress in human erythrocytes

AU May, James M.; Qu, Zhi-Chao; Xia, Li; Cobb, Charles E.

CS Departments of Medicine and Molecular Physiology and Biophysics, Vanderbilt University School of Medicine, Nashville, TN, 37232-6303, USA

SO American Journal of Physiology (2000), 279(6, Pt. 1), C1946-C1954

CODEN: AJPHAP; ISSN: 0002-9513

PB American Physiological Society

DT Journal

LA English

AB Nitric oxide, when released into the bloodstream, is quickly scavenged by Hb in erythrocytes or oxidized to nitrite. Nitrite can also enter erythrocytes and oxidize Hb. The goals of this work were to det. the mechanism of erythrocyte nitrite uptake and whether this uptake causes oxidant stress in these cells. Erythrocytes took up 0.8 mM nitrite with a half-time of 11 min. Nitrite uptake was sensitive to temp. and to the pH and ionic compn. of the medium but was not inhibited by the specific anion-exchange inhibitor

DIDS. About 25% of nitrite uptake occurred on the sodium-dependent phosphate transporter and the rest as diffusion of nitrous acid or other species across the plasma membrane. MetHb formation increased in proportion to the intracellular nitrite concn. Nitrite reacted with erythrocyte ~~ascorbate~~, but ~~ascorbate~~ loading of cells decreased nitrite-induced metHb formation only at high nitrite concns. In conclusion, nitrite rapidly enters erythrocytes and reacts with oxyHb but does not exert a strong oxidant stress on these cells.

IT 10102-43-9, Nitric oxide, biological studies
(nitrite uptake and metab. and oxidant stress in human erythrocytes in relation to)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

CC 13-6 (Mammalian Biochemistry)
IT 50-81-7, Ascorbic acid, biological studies 6730-29-6,
~~Ascorbate~~ radical, biological studies
(nitrite uptake and metab. and oxidant stress in human erythrocytes)
IT 10102-43-9, Nitric oxide, biological studies
(nitrite uptake and metab. and oxidant stress in human erythrocytes in relation to)
OSC.G 37 THERE ARE 37 CAPLUS RECORDS THAT CITE THIS RECORD (37 CITINGS)
RE.CNT 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L75 ANSWER 8 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 134:53243 HCA Full-text

TI An integrated nitric oxide sensor based on carbon fiber coated with selective membranes

AU Zhang, Xueji; Cardosa, Levis; Broderick, Mark; Fein, Harry; Lin, Jie
CS Department of Chemistry, World Precision Instruments, Inc., Sarasota, FL, 34240-9258, USA

SO Electroanalysis (2000), 12(14), 1113-1117
CODEN: ELANEU; ISSN: 1040-0397

PB Wiley-VCH Verlag GmbH

DT Journal

LA English

AB In vivo measurement of nitric oxide (NO) in a biol. matrix is very difficult because of its assumed low stability and fugacity, in addn. to the complexity of such matrix, limited space and vol. of biol. samples. Among different NO detection strategies, electrochem. NO sensors are still widely used by NO researchers. Though many kinds of NO sensors are com. available from World Precision Instruments, Inc. and other companies, the small NO sensors still are needed for the NO detection, esp. in single cell levels. In this article a NO-selective ultramicrosensor was developed as an easily

applicable tool for real time nitric oxide (NO) detection. The sensor consists of a 7 μ m carbon fiber working electrode coated with cation exchanger (Nafion), then covered with NO-selective gas permeable polymeric membranes, and Ag/AgCl micro-ref./counter electrode. Compared with other reported NO sensors, the sensor described herein offers several advantages: i) high selectivity against ascorbate (>104:1), dopamine (>103:1) and nitrite (104:1); ii) detection limit to low nanomolar concn.; iii) rapid, inexpensive and reproducible fabrication; iv) wide linear calibration range from 10 nM to 5 μ M with R²=0.995; v) integrated ultramicrosensor eliminating the need of an external ref. electrode, accordingly, expts. in small vol. are possible with an integrated ultramicrosensor, even at single cell levels.

IT 10102-43-9, Nitric oxide, analysis
(nitric oxide detn. using amperometric electrode based on carbon fiber coated with selective membranes)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

CC 9-1 (Biochemical Methods)
IT Cation exchange membranes
(nitric oxide detn. using amperometric electrode based on carbon fiber coated with selective membranes)
IT 10102-43-9, Nitric oxide, analysis
(nitric oxide detn. using amperometric electrode based on carbon fiber coated with selective membranes)
OSC.G 44 THERE ARE 44 CAPLUS RECORDS THAT CITE THIS RECORD (45 CITINGS)
RE.CNT 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L75 ANSWER 9 OF 20 HCA COPYRIGHT 2010 ACS on STN
AN 132:277974 HCA Full-text
TI Bioactivities of a tumor necrosis-like factor released by chicken macrophages
AU Rautenschlein, Silke; Subramanian, Anuradha; Sharma, Jagdev M.
CS Department of Veterinary Patho Biology, University of Minnesota, St Paul, MN, 55108, USA
SO Developmental & Comparative Immunology (1999), 23(7-8), 629-640
CODEN: DCIMDQ; ISSN: 0145-305X
PB Elsevier Science Ltd.
DT Journal
LA English
AB To test for tumor necrosis-like factor (TNF) of chickens, supernatants of a lipopolysaccharide (LPS)-stimulated chicken macrophage cell line MQ-NCSU were analyzed. A sequence of ion-exchange and gel-permeation chromatog. was

utilized to isolate TNF-like activity from the culture supernatant. The peak of TNF-like cytotoxic activity corresponded to the fractions with a mol. wt. of 81 kDa or higher. Polyclonal anti-human TNF- α antiserum cross-reacted by Western blotting with a 17 kDa protein in the TNF-contg. fraction under denaturing conditions. This result indicated that chicken TNF-like factor in the biol. active form may be a protein multimer of monomers of about 17 kDa. The mol. wt. of these monomers is similar to the mol. wt. of mammalian TNF- α . Chicken TNF-like factor stimulated macrophages by inducing morphol. changes, enhancing Ia-expression, nitric oxide (NO) prodn. and by synergizing with interferon (IFN)- γ in the induction of NO release from macrophages. The biol. activities were not neutralized by anti-human TNF antiserum. These data suggest that LPS-stimulated chicken macrophages produced a functional homolog to mammalian TNF- α . This may be structurally quite different from the mammalian TNF mol. Other factors may have been co-purified with the chicken TNF-like factor having overlapping functions and mol. wt. However, co-purifn. of chemokines and interleukin-1, major macrophage derived factors, with the chicken TNF-like factor can be excluded based on the purifn. strategies.

IT 10102-43-9, Nitric oxide, biological studies
(bioactivities of a tumor necrosis-like factor released by chicken macrophages)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

CC 15-5 (Immunochemistry)
Section cross-reference(s): 12
IT 10102-43-9, Nitric oxide, biological studies
(bioactivities of a tumor necrosis-like factor released by chicken macrophages)
OSC.G 18 THERE ARE 18 CAPLUS RECORDS THAT CITE THIS RECORD (18 CITINGS)
RE.CNT 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L75 ANSWER 10 OF 20 HCA COPYRIGHT 2010 ACS on STN
AN 127:351684 HCA Full-text
OREF 127:68859a,68862a
TI Manufacture of platinum-carrying silica gel catalyst by ion exchange
IN Tsurumi, Kazunori
PA Tanaka Kikinzoku Kogyo K. K., Japan
SO Jpn. Kokai Tokkyo Koho, 5 pp.
CODEN: JKXXAF
DT Patent
LA Japanese
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 09276698	A	19971028	JP 1996-96589	19960418
PRAI	JP 1996-96589		19960418		
AB	The manufg. method involves a process of treating a SiO ₂ gel with a Pt(IV) ammine complex ion obtained by heat treating ammonium chloroplatinate(IV) with an aq. NH ₃ soln. and vaporizing the excess NH ₃ . The catalyst is useful for oxidizing SO ₂ , CO, NO, and NH ₃ and dehydrating or hydrating a hydrocarbon. The obtained catalyst has a high Pt sp. surface area and shows high catalytic properties with the lower content of Pt.				
IT	10102-43-9, Nitric oxide, reactions (manuf. of silica gel catalyst supporting platinum ammine complex by ion exchange)				
RN	10102-43-9 HCA				
CN	Nitrogen oxide (NO) (CA INDEX NAME)				

N=O

IC ICM B01J023-42
ICS B01J021-08; B01D053-86
CC 67-1 (Catalysis, Reaction Kinetics, and Inorganic Reaction Mechanisms)
IT 630-08-0, Carbon monoxide, reactions 7446-09-5, Sulfur dioxide, reactions 7664-41-7, Ammonia, reactions 10102-43-9, Nitric oxide, reactions 16919-58-7, Ammonium chloroplatinate(IV)
(manuf. of silica gel catalyst supporting platinum ammine complex by ion exchange)

L75 ANSWER 11 OF 20 HCA COPYRIGHT 2010 ACS on STN
AN 126:311373 HCA Full-text
OREF 126:60217a,60220a

TI Preparation of iron complexes containing 1,3-diamino-2-hydroxypropanetetra(acetic acid) and sulfite ligands as nitrogen monoxide adsorbents
IN Sato, Terubumi; Yamada, Takashi
PA Mizusawa Industrial Chemicals, Ltd., Japan
SO Jpn. Kokai Tokkyo Koho, 8 pp.
CODEN: JKXXAF
DT Patent
LA Japanese
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 09067385	A	19970311	JP 1995-223568	19950831
	JP 3589510	B2	20041117		
PRAI	JP 1995-223568		19950831		
AB	An iron complex contg. Fe(II) and/or Fe(III) ions and 1,3-diamino-2-hydroxypropanetetra(acetic acid) (DHPTA) and sulfite ligands represented by compn. formula $Mp[Fe_4(O)_2(SO_3)_2-n(SO_4)_n(dhpta)_2]$ [M = cation; dhpta = 1,3-				

diamino-2-hydroxypropanetetra(acetic acid); n = 0,1; p = no. satisfying mp = 2-10; m = valence no. of cation M] are prep'd. by reacting a water-sol. Fe(II) salt, 1,3-diamino-2-hydroxypropanetetra(acetic acid), and a water-sol. sulfite salt in a aq. solvent in nonoxidizing atm. followed by optional oxidn. An iron complex comprising an iron-complex anion contg. Fe(II) and/or Fe(III) ions and 1,3-diamino-2-hydroxypropanetetra(acetic acid) and sulfite ligands, preferably represented by compn. formula $[\text{Fe}_4(\text{O})_2(\text{SO}_3)_2-n(\text{SO}_4)_n(\text{dhpta})_2]^{k-}$ (n = 0,1; k = 2-10), which are bonded to an org. or inorg. anion exchanger, is prep'd. by reacting a water-sol. Fe(II) salt, 1,3-diamino-2-hydroxypropanetetra(acetic acid), and a water-sol. sulfite salt in a aq. solvent in nonoxidizing atm., mixing the product soln. with an org. or inorg. anion exchanger, sepg. the product, and optional oxidn. before or after mixing the anion exchanger. A nitrogen monoxide (NO)-adsorbent consisting of the above iron complex is claimed. These iron-complexes are useful as adsorbents for nitrogen oxide, in particular NO(g). FeSO₄·7H₂O 2.23, DHPTA 1.29, and NaHSO₃ 2.03 g were dissolved in 100 cm³ H₂O with stirring at 50° for 20 min under Ar to give an aq. soln. contg. an tetra-iron complex (2 mmol), which was stirred with .apprx.20 g methanol-washed Dowex 2X8 (Cl-form) under Ar. The Dowex resin was filtered off, washed with H₂O at 50° and then with MeOH, and air-dried to give a dry resin (.apprx.18 g). The dry resin-immobilized Fe(II) complex (7.8 g) was packed in a glass tube, to which was passed N contg. 902 ppm NO at 420 cm³/min. The NO removal ratio was initially 95% and after passing 95 L gas for 3.8 h, it became 0. A total accumulation of NO absorbed was 3.5 mmol.

IT 10102-43-9, Nitrogen monoxide, processes
 (prepn. of iron complexes contg.
 diaminohydroxypropanetetra(acetic acid) and sulfite ligands and
 anion exchanger-immobilized iron complexes as
 nitrogen monoxide adsorbents)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



IT 11138-20-8, Dowex 2X8
 (prepn. of iron complexes contg. diaminohydroxypropanetetra(acetic acid) and sulfite ligands and anion exchanger
 -immobilized iron complexes as nitrogen monoxide adsorbents)

RN 11138-20-8 HCA

CN Dowex 2X8 (CA INDEX NAME)

*** STRUCTURE DIAGRAM IS NOT AVAILABLE ***

IC ICM C07F015-02
 ICS B01J020-22

CC 78-7 (Inorganic Chemicals and Reactions)

IT Adsorbents
 (prepn. of iron complexes contg. diaminohydroxypropanetetra(acetic acid) and sulfite ligands and anion exchanger
 -immobilized iron complexes as nitrogen monoxide adsorbents)

IT **Anion exchangers**
 (prepn. of iron complexes contg. diaminohydroxypropanetetra(acetic acid) and sulfite ligands and **anion exchanger** -immobilized iron complexes as nitrogen monoxide adsorbents)

IT 1344-28-1, Aluminum oxide (Al₂O₃), reactions
 (Neobead DN 1A; prepn. of iron complexes contg. diaminohydroxypropanetetra(acetic acid) and sulfite ligands and **anion exchanger**-immobilized iron complexes as nitrogen monoxide adsorbents)

IT 10102-43-9, Nitrogen monoxide, processes
 (prepn. of iron complexes contg. diaminohydroxypropanetetra(acetic acid) and sulfite ligands and **anion exchanger**-immobilized iron complexes as nitrogen monoxide adsorbents)

IT 1314-23-4, Zirconia, reactions 3148-72-9,
 1,3-Diamino-2-hydroxypropanetetra(acetic acid) 7720-78-7, Iron(II) sulfate 7757-83-7, Sodium sulfite 11138-20-8,
 Dowex 2X8 13463-67-7, Titania, reactions
 (prepn. of iron complexes contg. diaminohydroxypropanetetra(acetic acid) and sulfite ligands and **anion exchanger** -immobilized iron complexes as nitrogen monoxide adsorbents)

IT 15438-31-0DP, Ferrous ion, complexes, preparation 20074-52-6DP,
 Ferric ion, complexes, preparation 189275-27-2P 189275-28-3P
 189275-29-4DP, exchanged on Dowex 2X8 189275-30-7DP,
 exchanged on Dowex 2X8
 (prepn. of iron complexes contg. diaminohydroxypropanetetra(acetic acid) and sulfite ligands and **anion exchanger** -immobilized iron complexes as nitrogen monoxide adsorbents)

L75 ANSWER 12 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 126:91507 HCA Full-text

OREF 126:17633a,17636a

TI Manufacture of nitric oxide and apparatus therefor

IN Hirose, Yasuo

PA Hitachi Ltd, Japan

SO Jpn. Kokai Tokkyo Koho, 16 pp.

 CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	JP 08290906	A	19961105	JP 1995-96507	19950421
PRAI	JP 1995-96507		19950421		

AB The process comprises adding water to the first aq. electrolyte contg. HNO₃ in the first anode chamber of a NO producing means having a first anoned chamber and a first cathode chamber, which are sepd. by an ion-exchange membrane, reducing HNO₃ in the second aq. electrolyte in the first cathode chamber by electrolysis to form NO, taking out NO from the NO producing means, feeding the second aq. electrolyte to the second anode chambers of a HNO₃ concg. means having multiple second anode chambers and second cathode chambers formed by alternately arranging cationic-exchange membranes and

anionic- exchange membranes, transferring water accompanied in H ion from the second anode chambers to the second cathode chambers by electrodialysis, and returning the second aq. electrolyte to the first cathode chamber. The process decreases electricity consumption. The app. is also claimed.

IT 10102-43-9P, Nitric oxide, preparation
(manuf. of nitric oxide by electrolysis of nitric acid and app.
therefor)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)

$\text{N}=\text{O}$

IC ICM C01B021-24
ICS G21C019-46
CC 49-3 (Industrial Inorganic Chemicals)
IT 10102-43-9P, Nitric oxide, preparation
(manuf. of nitric oxide by electrolysis of nitric acid and app.
therefor)

L75 ANSWER 13 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 122:152236 HCA Full-text

OREF 122:27969a,27972a

TI Cultured astrocytes release a factor that decreases endothelin-1 secretion by brain microvessel endothelial cells

AU Federici, C.; Camoin, L.; Creminon, C.; Chaverot, N.; Strosberg, A. D.; Couraud, P. O.

CS Lab. d'Immuno-Pharm. Mol., Univ. Paris VII, Gif-sur-Yvette, Fr.

SO Journal of Neurochemistry (1995), 64(3), 1008-15

CODEN: JONRA9; ISSN: 0022-3042

PB Lippincott-Raven

DT Journal

LA English

AB Endothelin-1 (ET-1), originally characterized as a potent vasoconstrictor peptide secreted by vascular endothelial cells, has now been described to possess a wide range of biol. activities within the cardiovascular system and in other organs. Brain microvessel endothelial cells, which, together with perivascular astrocytes, constitute the blood-brain barrier, have been shown to secrete ET-1, whereas specific ET-1 receptors are expressed on astrocytes. It is reported here that conditioned medium from primary cultures of mouse embryo astrocytes could significantly, and reversibly, attenuate the accumulation of both ET-1 and its precursor big ET-1 in the supernatant of rat brain microvessel endothelial cells by up to 59 and 76%, resp., as assessed by immunometric assay. This inhibitor of ET-1 prodn. was purified by gel-exclusion and ion-exchange chromatog. as a 280-Da iron-contg. mol., able to release nitrites upon degrdn. These results suggest that astrocytes, via release of an iron-nitrogen oxide complex, may be involved in a regulatory loop of ET-1 prodn. at the level of the blood-brain barrier.

IT 10102-43-9DP, Nitric oxide, iron complex

(astrocytes in release of factor to decrease endothelin-1 secretion
by brain microvessel endothelial cells)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

$\text{N}=\text{O}$

CC 2-10 (Mammalian Hormones)

IT 7439-89-6DP, Iron, complex nitrogen oxide 10102-43-9DP,

Nitric oxide, iron complex

(astrocytes in release of factor to decrease endothelin-1 secretion
by brain microvessel endothelial cells)

OSC.G 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

L75 ANSWER 14 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 120:365 HCA Full-text

OREF 120:87a,90a

TI Biochemical characterization of a membrane-bound enzyme responsible
for **generating nitric oxide** from
nitroglycerin in vascular smooth muscle cells

AU Seth, Prem; Fung, Ho Leung

CS Sch. Pharm., State Univ. New York, Buffalo, NY, 14260, USA

SO Biochemical Pharmacology (1993), 46(8), 1481-6

CODEN: BCPA6; ISSN: 0006-2952

DT Journal

LA English

AB A membrane-bound enzyme responsible for **generating nitric oxide** (NO) from
nitroglycerin (NTG) in vascular smooth muscle cells has been characterized.
The enzyme could be solubilized from vascular microsomes by several
detergents, the most effective of which was 3-[(3-
cholamidopropyl)dimethylamino]-1- propanesulfonate (CHAPS). A partially
purified enzyme prepn. was obtained with CHAPS-solubilized vascular
microsomes that were processed sequentially through an **ion exchange** column
and a **gel** filtration column. The activity of this partially purified enzyme
showed a dependence on substrate concn., protein concn. and the duration of
incubation. Enzyme activity was enhanced 2.7- to 4.2-fold by several thiols
such as cysteine, N-acetylcysteine, reduced glutathione, and dithiothreitol.
On the other hand, N-ethylmaleimide, iodoacetic acid, p-chloromercuric
benzoic acid and 1-chloro-2,4-dinitrobenzene, reagents known to bind with
the free sulfhydryl groups, inactivated the NO-generating activity from NTG.
The enzyme activity could be reversibly bound to an organomercurial column.
These results suggested the presence of a free thiol group in the enzyme and
that this thiol group was required for enzyme activity. The partially
purified enzyme was active in the presence of 0.1% sodium dodecyl sulfate
(SDS). The enzyme was purified to near homogeneity using several sequential
chromatog. steps including DEAE-Sephacel, Biogel A 1.5 m, hydroxylapatite
and organomercurial columns, resulting in an increase in enzyme activity of
about 94-fold. The subunit of this enzyme, as identified on an SDS-treated
electrophoresis gel, had an apparent mol. size of 58 kDa.

IT 10102-43-9, Nitric oxide, biological studies
(nitroglycerin vasodilation mediation by formation of, by
membrane-bound enzyme)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)

$\text{N}=\text{O}$

CC 1-8 (Pharmacology)
Section cross-reference(s): 7
IT Thiols, biological studies
(nitric oxide generation by
membrane-bound enzyme dependence on, nitrovasodilators in relation
to)
IT Vasodilators
(nitro-, nitric oxide generation by,
membrane-bound enzyme mediation of)
IT 10102-43-9, Nitric oxide, biological studies
(nitroglycerin vasodilation mediation by formation of, by
membrane-bound enzyme)
IT 125978-95-2, Nitric oxide synthase
(of coronary microsomes, nitrovasodilator action mediation by
nitric oxide generation by)
IT 55-63-0P, Nitroglycerin
(vasodilation by, nitric oxide
generation by membrane-bound enzyme in)
OSC.G 25 THERE ARE 25 CAPLUS RECORDS THAT CITE THIS RECORD (25
CITINGS)

L75 ANSWER 15 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 118:216318 HCA Full-text

OREF 118:37245a,37248a

TI Method for removing cations and anions from an engine coolant liquid

IN Shubert, David C.; Myers, Galen R.; Richardson, Robert C.

PA BG Products, Inc., USA

SO U.S., 33 pp.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	US 5174902	A	19921229	US 1990-485939	19900227
PRAI	US 1990-485939		19900227		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A method and app. for removing particulates, hydrocarbons such as oil,
cations, and anions including NO₂- from a liq. such as automobile engine
coolant uses activated C filters and ion exchange beds. The app. has ≥1

filter for removing particulates and hydrocarbons; a strong acid cation exchange bed in the H form; a strong base anion exchange bed in the OH- form for removing anions; and a separator for sepg. gas contg. N, such as NO and/or NO₂, that is produced in the cation exchange bed and/or the anion exchange bed.

IT 10102-43-9P, Nitric oxide, preparation
(formation and removal of, from nitrites, in
anion exchange treatment of automobile engine
coolants)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

IC ICM C02F009-00
INCL 210662000
CC 51-11 (Fossil Fuels, Derivatives, and Related Products)
ST cation anion removal engine coolant liq; antifreeze impurity removal;
nitrite removal engine coolant liq
IT Antifreeze substances
(cations and anions in, removal of, by ion
exchange and adsorption, method and app. for)
IT Nitrites
(removal of, from automobile engine coolants, by anion
exchange, method and app. for)
IT Cooling agents
(liq., cations and anions in, removal of, by ion
exchange and adsorption, method and app. for)
IT 7782-77-6P, Nitrous acid 10102-43-9P, Nitric oxide,
preparation 10102-44-0P, Nitrogen dioxide, preparation
(formation and removal of, from nitrites, in
anion exchange treatment of automobile engine
coolants)
OSC.G 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)
RE.CNT 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT


L75 ANSWER 16 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 115:5803 HCA Full-text

OREF 115:1139a,1142a

TI Comparison of properties of nitric oxide and endothelium-derived
relaxing factor: some cautionary findings
AU Furchgott, R. F.; Khan, M. T.; Jothianandan, D.
CS Health Sci. Cent., SUNY, Brooklyn, NY, 11203, USA
SO Endothelium-Deriv. Relaxing Factors, Int. Symp. Endothelium-Deriv.
Vasoact. Factors, 1st (1990), Meeting Date 1989, 8-21.
Editor(s): Rubanyi, Gabor M.; Vanhoutte, Paul M. Publisher: Karger,
Basel, Switz.
CODEN: 57ATAZ

DT Conference
LA English
AB The differential sensitivity of smooth muscles to endothelium-derived relaxing factor (EDRF) and NO, the ability of an ~~anion- exchange~~ resin (1°, 2°-amino (NH₂/NH)) to remove the vascular relaxing activity of both EDRF and NO, and the appearance of NO₂⁻ as a major oxidn. product of NO, whether the oxidant is O₂ or O₂⁻ are demonstrated, and considerations on making and biol. testing of solns. of NO are discussed. The relevance of the findings of these expts. to the identity of EDRF and NO is discussed.
IT 14797-65-0, Nitrite, biological studies
(as nitric oxide oxidn. product)
RN 14797-65-0 HCA
CN Nitrite (8CI, 9CI) (CA INDEX NAME)



IT 10102-43-9P, Nitric oxide, biological studies
(endothelium-derived relaxing factor identity with, nitric oxide
prepn. and testing in relation to)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)



CC 13-6 (Mammalian Biochemistry)
IT 14797-65-0, Nitrite, biological studies
(as nitric oxide oxidn. product)
IT 10102-43-9P, Nitric oxide, biological studies
(endothelium-derived relaxing factor identity with, nitric oxide
prepn. and testing in relation to)
OSC.G 14 THERE ARE 14 CAPLUS RECORDS THAT CITE THIS RECORD (14
CITINGS)

L75 ANSWER 17 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 96:71455 HCA Full-text

OREF 96:11737a,11740a

TI Determination of total N-nitroso content in cutting fluids

AU Cox, Robert D.; Frank, Clyde W.

CS Dep. Chem., Univ. Iowa, Iowa City, IA, 52242, USA

SO Analytical Chemistry (1982), 54(3), 557-9

CODEN: ANCHAM; ISSN: 0003-2700

DT Journal

LA English

AB A rapid detn. of the total N-nitroso content of cutting fluids involves initial removal of nitrite by ion exchange, iodide ion, or sulfanilamide [1116-54-7]. The nitrite-free sample is analyzed by denitrosation of N-

nitroso compds. to produce NO, which is detected via its gas-phase chemiluminescence reaction with O₃. The detection limit is 5 + 10⁻¹¹ mol on cutting fluid samples. Anal. time is 5-15 min.

IT 10102-43-9P, preparation
(formation of, in detn. of N-nitroso content of cutting fluids)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)



CC 51-8 (Fossil Fuels, Derivatives, and Related Products)
Section cross-reference(s): 80

IT 10102-43-9P, preparation
(formation of, in detn. of N-nitroso content of cutting fluids)

L75 ANSWER 18 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 65:16303 HCA Full-text

OREF 65:3038f-h,3039a

TI Sorption of carbon disulfide by anion-exchange resins

AU Tsaplina, L. A.; Davankov, A. B.

SO Zhurnal Prikladnoi Khimii (Sankt-Peterburg, Russian Federation) (1966), 39(3), 608-11

CODEN: ZPKHAB; ISSN: 0044-4618

DT Journal

LA Russian

GI For diagram(s), see printed CA Issue.

AB The sorption of CS₂ from aq. solns. was investigated on the OH forms of the air-dried AN-1, MMG-1, N-O, EDE-10P, and SDTM anion-exchange resins as well as on AP-3 activated C under static conditions. The sorption capacity of the investigated systems to CS₂ is about the same. The greater the CS₂ concn. in the aq. soln., the greater the sorption. The sorption is accompanied by changes in resin color. These colors are derived from the interaction of CS₂ with the active resin groups, giving rise to the formation of a new type of xanthate, according to the reactions: These types of compds. are readily decompd. by 1% HCl solns. with the resin completely recovering its initial color and ion exchange capacity as detd. by expts. carried out by regenerating the resin. This was accomplished by alternate loading of the resin with 0.1N H₂SO₄ and HCl solns. The influence of the flow rate and temp. on the sorption process considered showed little effect on the 1st parameter, but increases in temp. resulted in a noticeable sorption increase. The anion-exchangers proved to be superior to the activated C as sorbents. Carrying out the sorption process at 40° brings about a 50% increase in sorption efficiency by the resins.

IT 10102-43-9P, N-O
(carbon disulfide adsorption by, xanthate formation in)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

CC 4 (Surface Chemistry and Colloids)
IT ~~Anion-exchanging~~ substances
(carbon disulfide adsorption by, xanthate formation in)
IT Adsorption
(of carbon disulfide, by ~~anion-exchange~~ resins,
xanthate formation in)
IT 75-15-0P, Carbon disulfide
(adsorption of, by ~~anion-exchange~~ resins,
xanthate formation in)
IT 9086-61-7P, AN 1 ~~10102-43-9P~~, N-O 11106-30-2P, EDE 10P
76483-21-1P, AP 3
(carbon disulfide adsorption by, xanthate formation in)
IT 151-01-9P, Xanthate
(formation of, in CS2 adsorption by ~~anion~~
~~exchange~~ resins)

L75 ANSWER 19 OF 20 HCA COPYRIGHT 2010 ACS on STN
AN 62:20775 HCA Full-text
OREF 62:3696f-g
TI Extraction of tungsten from nitric acid solutions
AU Yurkevich, Yu. N.; Sviridovskaya, R. M.
SO Sb. Tr. Vses. Nauchn.-Issled. Inst. Tverd. Splavov (1964),
(5), 245-9
From: Ref. Zh., Met. 1964, Abstr. No. 9G117.
DT Journal
LA Russian
AB The possibility of extg. WO3 from HNO3 solns. with the aid of the ~~anion~~
~~exchanger~~ N-O was studied. The total exchange capacity of the ~~exchanger~~ N-O
at an acidity of 5-50 g. HNO3/l. was not inferior to that of the ~~anion~~
~~exchanger~~ EDE-10P in HCl solns., and was 170-80 kg. WO3/ton ion exchanger.
W was regenerated with 10% NaOH. WO3 can be obtained from the regenerated
product by existing procedures.
IT ~~10102-43-9P~~, N-O
(in tungsten extn. from HNO3 solns.)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

CC 18 (Extractive Metallurgy)
IT ~~Anion exchange~~
(in tungsten extn. from HNO3 solns.)
IT ~~10102-43-9P~~, N-O
(in tungsten extn. from HNO3 solns.)

IT 7440-33-7, Tungsten
(process metallurgy of, from nitric acid soln. by anion
exchange)

L75 ANSWER 20 OF 20 HCA COPYRIGHT 2010 ACS on STN

AN 62:19176 HCA Full-text

OREF 62:3443e-f

TI Absorption of cations by various anion exchangers

AU Muromtseva, G. V.; Ol'shanova, K. M.; Saldadze, K. M.; Kopylova, V. D.

SO Issled. Svoistv Ionobmen. Materialov, Akad. Nauk SSSR, Inst. Fiz.
Khim. (1964) 108-14

DT Journal

LA Russian

AB The influence of structure of various Soviet anion exchange resins, and their salt form, pretreatment, and temp. on the sorption capacity for Cu++ was studied. The anion exchangers in Cl form were brought in contact with 0.1N CuCl2 of pH 3.8. Monofunctional anion exchangers of polymn. type do not sorb Cu++, whereas those obtained by polycondensation form complexes with Cu++. Cu++ is uniformly distributed inside the beads. For complex formation, the presence of primary and secondary amine groups is necessary, and their complexing capacity is increased by tertiary amine and OH groups. The complexes are split by acids and not by NH3. With anion exchangers in OH form, sparingly sol. Cu(OH)2 or basic salts are formed mainly on the bead surface. Condensation-type anion exchangers reduce Ag+ to its metal form. With increasing temp., the sorption capacity increases. Pretreatment of com. anion exchangers has no effect.

IT 10102-43-9P, N-O
(anion exchange capacity of, complex formation
and)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

$\text{N}=\text{O}$

CC 4 (Surface Chemistry and Colloids)

IT Anion exchange
(capacity of)

IT Amino group
(in anion-exchanging resins, complex formation
and)

IT Hydroxyl group
(on anion-exchanging resins, complex formation
and)

IT 7440-50-8P, Copper
(and salts, basic, formation of, in anion
exchange of Cu)

IT 9064-43-1P, AN 2FG 9086-61-7P, AN 1 10102-43-9P, N-O
11106-30-2P, EDE 10P 11111-77-6P, AV 16 11138-00-4P, AN-15
12640-33-4P, AN-31 30176-85-3P, Phenol,

2,2'-thiobis[tert-butyl-4-chloro- 37380-46-4P, AN-20 39454-56-3P,
AV 18 56939-65-2P, AN 17 61642-45-3P, AN-24

(anion exchange capacity of, complex formation
and)

IT 11106-27-7P, AV 17

(anion-exchange capacity of, complex formation
and)

IT 20427-59-2P, Copper hydroxide, Cu(OH)2

(formation of, in anion exchange of Cu)

IT 7440-22-4, Silver

(redn. of, by anion-exchange resins)

=> D L78 1-14 BIB ABS HITSTR HITIND

L78 ANSWER 1 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 139:207973 HCA Full-text

TI Estrone and estradiol mediate vascular function by different
mechanisms

AU Massheimer, V.; Polini, N.; Benozzi, S.; Alvarez, C.; Selles, J.

CS Catedra de Analisis Clinicos II, Departamento de Biologia, Bioquimica
y Farmacia. Universidad Nacional del Sur, Bahia Blanca, B8000ICN,
Argent.

SO Revista Argentina de Endocrinologia y Metabolismo (2003),
40(1), 3-12

CODEN: RAEMA7; ISSN: 0326-4610

PB Sociedad Argentina de Endocrinologia y Metabolismo

DT Journal

LA Spanish

AB Postmenopausal women have an increased risk for cardiovascular disease which
is assocd. to the lost of the vascular protective action of estradiol. In
this period circulating estradiol (E2) levels are considerably low, but
estrone (E1) levels remain high due to its peripheral synthesis. The
endothelium produces different active metabolites, such as NO and
eicosanoids (thromboxane, prostaglandins and prostacycline), which regulate
arterial vasomotor properties and platelet aggregation. Previously the
authors demonstrated that rat aortic tissue treated with physiol. concns. of
estradiol and progesterone for 1-5 min inhibits platelet aggregation
mediated by NOS activation. The authors also reported progesterone rapid
action on aortic cyclooxygenase. The aim of the present study was to
compare the mechanism involved in the rat aortic rapid response to E1 or E2.
Rat aortic strips (RAS) with intact endothelium, were treated in vitro for
1-5 min with physiol. concns. of E2 or E1. Platelet aggregation (PA)
induced by 10 μ M ADP was measured in a platelet-rich plasma (PRP) which was
incubated with RAS and treated with the hormones. NO prodn. was measured by
conversion of 3H-arginine to 3H-citruline reaction. 3H-citruline was detd.
by ion exchange chromatog. using a Dowex AG-50WX8 column. Eicosanoids
prodn. was measured by TLC using 3H-arachidonic acid as precursor. The
increase in NO prodn. induced by 1 nM E1 treatment was abolished by the
presence of L-NAME in the incubation media, confirming that NOS is activated
in aortic tissue in response to E1 as has been demonstrated for E2. Calcium

requirement for aorta NOS rapid activation by E1 and E2 treatment was studied. The presence of 0.5 mM EGTA in the incubation medium abolished the increase in NO prodn. induced by 1 nM E2, whereas the tissue response to 1 nM E1 was not affected by the calcium chelator, implying that the aorta response to E1 treatment does not require extracellular Ca²⁺. Both E2 and E1 treatment inhibited PA, but the effect elicited by E1 is less potent compared with E2. The eicosanoid signal transduction pathway is involved in RAS rapid response to E2 or E1. The authors' results show that both hormones increase PGI₂ release by aortic tissue, with higher stimulus induced by E1. Thromboxane (Tx) prodn. was stimulated only by E1. Considering the potent effect of Tx on platelet aggregation, the authors detd. the effect of E1 treatment on platelet aggregation in the presence of the cyclooxygenase (COX) inhibitor indomethacine. The authors found that under this exptl. condition E1 treatment produced an inhibition of platelet aggregation equiv. to that elicited by E2. These results suggest that E1 and E2 modulate rat aortic NOS and COX activity by different mechanisms.

IT 10102-43-9, Nitric oxide, biological studies
 (estradiol and estrone effects on eicosanoid and ~~nitric~~
~~oxide formation~~ and platelet aggregation in rat
 aorta mediation by different mechanisms)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

CC 2-4 (Mammalian Hormones)

IT Artery
 (aorta; estradiol and estrone effects on eicosanoid and
~~nitric oxide formation~~ and platelet
 aggregation in rat aorta mediation by different mechanisms)

IT Platelet aggregation
 Platelet aggregation
 (estradiol and estrone effects on eicosanoid and ~~nitric~~
~~oxide formation~~ and platelet aggregation in rat
 aorta mediation by different mechanisms)

IT 50-28-2, Estradiol, biological studies 53-16-7, Estrone, biological
 studies 10102-43-9, Nitric oxide, biological studies
 35121-78-9, PGI₂ 39391-18-9, Cyclooxygenase 125978-95-2, Nitric
 oxide synthase
 (estradiol and estrone effects on eicosanoid and ~~nitric~~
~~oxide formation~~ and platelet aggregation in rat
 aorta mediation by different mechanisms)

OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

RE.CNT 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L78 ANSWER 2 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 138:149777 HCA Full-text

TI Evaluation of methods for the extraction of ~~nitrite~~ and

nitrate in biological fluids employing high-performance anion-exchange liquid chromatography for their determination

AU Smith, Christopher C. T.; Stanyer, Lee; Betteridge, D. John

CS Department of Medicine, The Middlesex Hospital, Royal Free and University College Medical School, London, W1N 8AA, UK

SO Journal of Chromatography, B: Analytical Technologies in the Biomedical and Life Sciences (2002), 779(2), 201-209
CODEN: JCBAAI; ISSN: 1570-0232

PB Elsevier Science B.V.

DT Journal

LA English

AB Measurements of NO₂⁻ and NO₃⁻ in biol. fluids are proposed as indexes of cellular NO prodn. Detn. of NO₂⁻ and NO₃⁻ in std. solns. is not difficult, however, detns. which reflect accurately cellular NO synthesis represent a considerable anal. challenge. Problems are often encountered arising from background NO₂⁻/NO₃⁻ contamination in exptl. solns. and lab. hardware, and with methods for sample extn. We investigated potential procedures for the extn. and detn. of NO₂⁻ and NO₃⁻ in biol. samples. Consequently, a protocol was devised which yielded acceptable results regarding extn. efficiency, assay reproducibility, sample throughput and contaminant minimization. It entailed rigorous washing of all equipment with water of low NO₂⁻ and NO₃⁻ content, sample deproteinization by centrifugal ultrafiltration through a 3K filter, and anal. by high-performance anion-exchange liq. chromatog. with UV detection. Retention times for NO₂⁻ and NO₃⁻ in stds. and plasma were 4.4 and 5.6 min, resp. Assay linearity for stds. ranged between 31 nM and 1 mM. The limit of detection for NO₂⁻ and NO₃⁻ in stds. was 3 pmol. Recoveries of NO₂⁻ and NO₃⁻ from spiked plasma (1-100 µM KNO₂/KNO₃) and from extd. stds. (1-250 µM) were .apprx.100%. Intra-assay and inter-assay RSDs for NO₂⁻ and NO₃⁻ in spiked and unspiked plasma were ≤10.6%. Assays on washed platelet supernatants demonstrated collagen-induced platelet generation of NO products and anal. of murine and rat cardiac perfusates was achieved. Our procedure may be suitable for routine detn. of NO₂⁻ and NO₃⁻ in various biol. fluids, e.g., plasma.

IT 10102-43-9, Nitric oxide, analysis
(extn. of nitrite and nitrate in biol. fluids employing high-performance anion-exchange liq. chromatog. for their detn.)

RN 10102-43-9 HCA


CN Nitrogen oxide (NO) (CA INDEX NAME)

N=O

IT 14797-55-8P, Nitrate, analysis 14797-65-0P, Nitrite, analysis
(extn. of nitrite and nitrate in biol. fluids employing high-performance anion-exchange liq. chromatog. for their detn.)


RN 14797-55-8 HCA

CN Nitrate (6CI, 7CI, 8CI, 9CI) (CA INDEX NAME)



RN 14797-65-0 HCA

CN Nitrite (8CI, 9CI) (CA INDEX NAME)



CC 9-9 (Biochemical Methods)

ST nitrate ~~nitrite~~ extn blood HPLC

IT ~~Anion exchange~~ HPLC

Blood analysis

Extraction

Platelet (blood)

(extn. of ~~nitrite~~ and nitrate in biol. fluids employing
high-performance ~~anion-exchange~~ liq. chromatog.
for their detn.)

IT 10102-43-9, Nitric oxide, analysis

(extn. of ~~nitrite~~ and nitrate in biol. fluids employing
high-performance ~~anion-exchange~~ liq. chromatog.
for their detn.)

IT 14797-55-8P, Nitrate, analysis 14797-65-0P,
~~Nitrite~~, analysis

(extn. of ~~nitrite~~ and nitrate in biol. fluids employing
high-performance ~~anion-exchange~~ liq. chromatog.
for their detn.)

OSC.G 20 THERE ARE 20 CAPLUS RECORDS THAT CITE THIS RECORD (20
CITINGS)

RE.CNT 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L78 ANSWER 3 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 122:299108 HCA Full-text

OREF 122:54329a,54332a

TI Polymer-bound nitric oxide/nucleophile adduct compositions for
treatment of biological disorders

IN Keefer, Larry K.; Hrabie, Joseph A.

PA United States Dept. of Health and Human Services, USA

SO U.S., 13 pp.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 11

PATENT NO.

KIND

DATE

APPLICATION NO.

DATE

PI	US 5405919	A	19950411	US 1992-935565	19920824
	US 5525357	A	19960611	US 1993-121169	19930914
	US 5650447	A	19970722	US 1994-214372	19940317
	US 5632981	A	19970527	US 1994-344157	19941122
	US 5676963	A	19971014	US 1995-417917	19950406
	US 5718892	A	19980217	US 1995-417913	19950406
	US 5691423	A	19971125	US 1995-419424	19950410
	US 5910316	A	19990608	US 1995-419044	19950410
	US 6110453	A	20000829	US 1998-13349	19980126
	US 6290981	B1	20010918	US 1999-289570	19990409
	US 6379660	B1	20020430	US 2000-666668	20000920
	US 20020119115	A1	20020829	US 2002-41200	20020108
	US 7425218	B2	20080916		
PRAI	US 1992-935565	A2	19920824		
	US 1993-121169	A2	19930914		
	US 1994-344157	A3	19941122		
	US 1995-417913	A3	19950406		
	US 1995-419044	A3	19950410		
	US 1997-837812	A1	19970422		
	US 2000-666668	A1	20000920		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A polymeric compn. capable of releasing nitric oxide comprises polymer and a nitric oxide-releasing functional group bound to the polymer for treatment of biol. disorders. The compns. can be used as and/or incorporated into implants, injectables, condoms, prosthesis coatings, patches, and the like for use in a wide variety of medical applications (no data). For example, poly(aminostyrene) in acetonitrile was placed under 5 atm nitric oxide to give a cream-colored polymer of which one-third of the amino side chains became attached to N2O2 groups.

IT ~~10102-43-9DP~~, Nitric oxide, polymer conjugates
(polymer-bound nitric oxide/nucleophile adducts for treating biol. disorders)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



IT ~~9002-88-4D~~, Polyethylene, nitric oxide conjugates
(polymer-bound nitric oxide/nucleophile adducts for treating biol. disorders)

RN 9002-88-4 HCA

CN Ethene, homopolymer (CA INDEX NAME)

CM 1

CRN 74-85-1

CMF C2 H4

H₂C=CH₂

IC ICM C08K005-22
ICS A01N033-26; A61K031-785; C08F008-30
INCL 525377000
CC 63-6 (Pharmaceuticals)
IT 9002-98-6DP, Polyethylenimine, nitric oxide conjugates 9060-90-6DP,
Poly(aminostyrene), nitric oxide conjugates ~~10102-43-9DP~~,
Nitric oxide, polymer conjugates 26780-50-7DP, Glycolide-lactide
copolymer, nitric oxide conjugates
(polymer-bound nitric oxide/nucleophile adducts for treating biol.
disorders)
IT 9002-84-0D, Polytetrafluoroethylene, nitric oxide conjugates
9002-86-2D, Polyvinyl chloride, nitric oxide conjugates
~~9002-88-4D~~, Polyethylene, nitric oxide conjugates
9003-07-0D, Polypropylene, nitric oxide conjugates 9003-53-6D,
Polystyrene, nitric oxide conjugates 24937-79-9D, Polyvinylidene
difluoride, nitric oxide conjugates
(polymer-bound nitric oxide/nucleophile adducts for treating biol.
disorders)
OSC.G 54 THERE ARE 54 CAPLUS RECORDS THAT CITE THIS RECORD (61
CITINGS)
RE.CNT 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L78 ANSWER 4 OF 14 HCA COPYRIGHT 2010 ACS on STN
AN 121:136789 HCA Full-text
OREF 121:24707a,24710a
TI Measurement of intraparticle effective diffusion coefficient of NO in
metal ~~ion-exchanged~~ zeolites by analysis of
breakthrough curves
AU Zhang, Wen Xiang; Yahiro, Hidenori; Izumi, Jun; Iwamoto, Masakazu
CS Catalysis Res. Cent., Hokkaido Univ., Sapporo, 060, Japan
SO Nippon Kagaku Kaishi (1994), (8), 748-51
CODEN: NKAKB8; ISSN: 0369-4577
DT Journal
LA Japanese
AB Breakthrough curves of NO adsorption on various metal ~~ion-exchanged~~
zeolites have been employed to evaluate the intraparticle effective
diffusion coeff. (Di). The Di was 0.7 + 10⁻³⁻²⁹ + 10⁻³ cm²/s and was
charged with zeolite structures, metal ~~ions exchanged~~, and adsorption temp.
On MFI zeolite, Di was dependent on the radius of metal ion, and a max. Di
was obsd. around 0.09 nm of the radius. With Cu-ZSM-5 and Agmordenite, the
max. Di was obsd. around 250 K, while the Di of Comordenite was not varied
with the adsorption temp.
IT ~~10102-43-9P~~, Nitrogen oxide (NO), preparation
(adsorption of, on metal ~~ion-exchanged~~
zeolites)
RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

$\text{N}=\text{O}$

CC 48-1 (Unit Operations and Processes)
ST diffusion nitrogen oxide ion exchanged zeolite;
metal ion exchanged zeolite adsorption;
intraparticle effective diffusion nitrogen oxide; breakthrough curve
adsorption nitrogen oxide
IT Adsorption
(of nitrogen oxide, in metal ion-exchanged
zeolites, measurement of intraparticle effective diffusion in)
IT 10102-43-9P, Nitrogen oxide (NO), preparation
(adsorption of, on metal ion-exchanged
zeolites)
OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L78 ANSWER 5 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 117:99411 HCA Full-text

OREF 117:17131a,17134a

TI Nitrogen oxides generation method for recovered nitric acid by
electrolysis. An action plan for reduction of low-level-liquid-waste
in processing plant

AU Suzuki, Kaunori

CS Oarai Nucl. Res. Cent., JGC Corp., Japan

SO Kyoto Daigaku Genshiro Jikkensho, [Tech. Rep.] (1991),

KURRI-TR-361, 19-26

CODEN: KDGHDH; ISSN: 0287-9808

DT Report

LA Japanese

AB A specified concn. HNO_3 was fed to an electrolytic cell and qual. and quant.
anal. of the gas formed were carried out. The main test parameters were
 HNO_3 concn. (1-12 mol/L), electrode material (Pt, graphite), c.d. (0.01-0.05
A/cm²), presence or absence of diaphragm (cationic exchange membrane) and
flow rate of HNO_3 in the electrolytic cell. The detns. of NO and NO₂ were
carried out by using a NO_x analyzer. The total NO_x was detd. by ozone
oxidn./alkali absorption/neutralization titrn., and H₂ and N₂O were detd. by
gas chromatog. The current efficiency (%) for the formation of NO_x was
calcd. by the equation: [amt. of NO₂ (mol/h) produced 26.8 (A-h/mol) + amt.
of NO (mol/h) produced 80.4 (A-h/mol) + amt. of NO (mol/h) produced 80.4 (A-
h/mol)] + 100/electricity (A) supplied. At high HNO_3 concn. a mixt. of NO
and NO₂ was produced. At medium HNO_3 concn. the main product was H₂ gas
when the HNO_3 concn. was ≤ 6 mol/L and Pt cathode was used whereas a mixt. of
NO and N₂O was produced when the HNO_3 concn. was 2-4 mol/L and graphite
electrode was used, however when the HNO_3 concn. was ≤ 1 mol/L H₂ was
produced. The current efficiency for high concn. HNO_3 electrolysis was $\geq 90\%$
so NO_x was formed effectively. When a diaphragm-contg. electrolytic cell
was used the prodn. efficiency of NO_x did not drop even when the flow rate

was small and the prodn. efficiency was $\geq 90\%$ whereas in an electrolytic cell without a diaphragm, the same efficiency as diaphragm cell was not obtained unless the flow rate (linear velocity) was large.

IT ~~10102-43-9P~~, Nitrogen monoxide, preparation
(prodn. of, from recovered nitric acid by electrolysis, radioactive waste redn. at reprocessing facility in relation to)
RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)



CC 71-11 (Nuclear Technology)
Section cross-reference(s): 72

IT ~~Cation exchangers~~
(membranes, for electrolytic cells for nitric acid recovery, radioactive waste redn. issues in relation to)
IT 1333-74-0P, Hydrogen, preparation ~~10102-43-9P~~, Nitrogen monoxide, preparation 10102-44-0P, Nitrogen dioxide, preparation 11104-93-1P, Nitrogen oxide, preparation
(prodn. of, from recovered nitric acid by electrolysis, radioactive waste redn. at reprocessing facility in relation to)

L78 ANSWER 6 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 115:60588 HCA Full-text

OREF 115:10279a,10282a

TI Approach to De-NO_x-ing photocatalysis. Photocatalytic decomposition of NO on Cu⁺/SiO₂ catalyst prepared via ~~ion-exchange~~ method

AU Anpo, Masakazu; Nomura, Takaiki; Kitao, Teijiro; Giamello, Elio; Che, Michel; Fox, Marye Anne

CS Coll. Eng., Univ. Osaka Prefect., Sakai, 591, Japan

SO Chemistry Letters (~~1991~~), (5), 889-92

CODEN: CMLTAG; ISSN: 0366-7022

DT Journal

LA English

AB Cu²⁺ ions supported onto SiO₂ (Cu²⁺/SiO₂) prepd. by an ~~ion-exchange~~ method are reduced to Cu⁺ ions when the Cu²⁺/SiO₂ sample is evacuated >573 K. Cu⁺/SiO₂ catalyst decomp. NO photocatalytically and stoichiometrically at 275 K. The excited state of the Cu⁺ ions plays a significant role in the photocatalytic decompn. of NO on the Cu⁺/SiO₂ catalyst.

IT ~~10102-43-9P~~, Nitrogen monoxide, reactions
(photocatalytic decompn. of, on copper ion(1+)-silica catalyst prepd. by ~~ion-exchange~~ method)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



CC 74-1 (Radiation Chemistry, Photochemistry, and Photographic and Other
Reprographic Processes)
ST photocatalyst copper ion exchange silica; nitrogen
oxide decompn photocatalyst copper ion; photodecompn nitrogen oxide
copper ion catalyst
IT Photolysis catalysts
(copper ion(1+)/silica, for nitrogen monoxide decompn., prepd. by
ion-exchange method)
IT Photolysis
(of nitrogen monoxide on copper ion/silica catalyst prepd. by
ion-exchange method)
IT 7631-86-9P, Silica, uses and miscellaneous
(photocatalyst contg. copper(1+) on, prepd. by ion-
exchange, for decompn. of nitrogen monoxide)
IT 17493-86-6P, Copper ion(1+), uses and miscellaneous
(photocatalyst from silicon dioxide and, prepd. by ion-
exchange, for nitrogen monoxide decompn.)
IT 10102-43-9P, Nitrogen monoxide, reactions
(photocatalytic decompn. of, on copper ion(1+)-silica catalyst
prepd. by ion-exchange method)
OSC.G 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD (9 CITINGS)

L78 ANSWER 7 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 114:48639 HCA Full-text

OREF 114:8317a,8320a

TI Rates and mechanisms of nitrogen dioxide removal from indoor air by
residential materials

AU Spicer, C. W.; Coutant, R. W.; Ward, G. F.; Joseph, D. W.; Gaynor, A.
J.; Billick, I. H.

CS Battelle Mem. Inst., Columbus, OH, 43201, USA

SO Environment International (1989), 15(1-6), 643-54

CODEN: ENVIDV; ISSN: 0160-4120

DT Journal

LA English

AB The relative efficiencies for NO2 removal from indoor air by a large no. of
materials are presented with a discussion of the factors that influence the
removal process. The reaction with indoor surfaces represents a significant
sink for NO2, and that these reactions are effecting a considerable degree
of control over indoor NO2 levels. It seems likely that this control could
be enhanced by judicious selection of furnishings and construction
materials.

IT 9002-88-4, Polyethylene
(air pollution by, indoor, residential building and furnishing
materials in mitigation of)

RN 9002-88-4 HCA

CN Ethene, homopolymer (CA INDEX NAME)

CM 1

CRN 74-85-1

CMF C2 H4

$\text{H}_2\text{C}=\text{CH}_2$

IT 10102-43-9P, Nitric oxide, preparation
(formation of, in nitrogen dioxide removal from indoor air by
residential building and furnishing materials)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

$\text{N}=\text{O}$

CC 59-2 (Air Pollution and Industrial Hygiene)

Section cross-reference(s): 40, 58

IT 9002-88-4, Polyethylene 10102-44-0, Nitrogen dioxide,
biological studies
(air pollution by, indoor, residential building and furnishing
materials in mitigation of)

IT 10102-43-9P, Nitric oxide, preparation
(formation of, in nitrogen dioxide removal from indoor air by
residential building and furnishing materials)

OSC.G 7 THERE ARE 7 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)

L78 ANSWER 8 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 110:14873 HCA Full-text

OREF 110:2487a,2490a

TI Experiments on acid digestion and gas-purification processes

AU Matteman, J. L.; De Niet, J.; Boekschoten, H. J. C.

CS Res. Dev. Div., N. V. Kema, Arnhem, 6800 ET, Neth.

SO Kema Scientific & Technical Reports (1988), 6(6), 133-44

CODEN: KESRED; ISSN: 0167-8590

DT Journal

LA English

AB Research carried out on acid digestion and gas purifn. started with a
selection of H_2SO_4 and as the chems. to be used. The design parameters for
low-level waste were then detd. in a pilot plant. The digestion of the
waste in the pilot plant resulted in the formation of SO_2 and NO_x . These
comps. were converted back to the corresponding acids in a gas-purifn.
system consisting of a series of contact columns. Reconversion of H_2SO_4
could be done in a relatively small column in which the SO_2 formed during
the digestion was oxidized by HNO_3 . The HNO_3 was recovered by absorbing NO_x
into water in three columns operating at room temp. Both air and
(preferably) H_2O_2 were successfully used as oxidants during absorption.
Experience indicated that an acid-digestion plant can be run in a safe and
reliable way in spite of the fact that aggressive chems. have to be used.
The burdening of the environment with NO_x or SO_2 is limited. The pilot

plant could be run by 1 person, owing to the installation of a process computer.

IT 9002-88-4, Polyethylene
(acid digestion of radioactive low-level waste contg., gas purifn.
in relation to)


RN 9002-88-4 HCA

CN Ethene, homopolymer (CA INDEX NAME)

CM 1

CRN 74-85-1

CMF C2 H4



IT 10102-43-9P, Nitrogen monoxide, preparation
(formation of, in acid digestion of radioactive low-level waste
contg. org. materials)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



CC 71-11 (Nuclear Technology)

IT 9002-86-2, Polyvinyl chloride 9002-88-4, Polyethylene
9004-34-6, Cellulose, reactions
(acid digestion of radioactive low-level waste contg., gas purifn.
in relation to)

IT 124-38-9P, Carbon dioxide, preparation 630-08-0P, Carbon monoxide,
preparation 7727-37-9P, Nitrogen, preparation 7732-18-5P, Water,
vapor 7782-44-7P, Oxygen, preparation 10102-43-9P,
Nitrogen monoxide, preparation 10102-44-0P, Nitrogen dioxide,
preparation
(formation of, in acid digestion of radioactive low-level waste
contg. org. materials)

L78 ANSWER 9 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 107:239285 HCA Full-text

OREF 107:38439a,38442a

TI Recovery of hydroxylamine or its salts from wastewaters

IN Fuchs, Hugo; Thomas, Erwin; Weiss, Franz Josef; Ritz, Josef

PA BASF A.-G., Fed. Rep. Ger.

SO Ger. Offen., 3 pp.

CODEN: GWXXBX

DT Patent

LA German

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	DE 3607998	A1	19870917	DE 1986-3607998	19860311
	US 4725360	A	19880216	US 1987-22875	19870306
	EP 236993	A2	19870916	EP 1987-103283	19870307
	EP 236993	A3	19880504		
	JP 62213893	A	19870919	JP 1987-52250	19870309
PRAI	DE 1986-3607998	A	19860311		

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The title process is carried out by passing the wastewaters over a strongly acidic **ion-exchange** resin, and then contacting the **ion-exchange** resin with aq., 5-15-wt.% H2SO4 to obtain an aq. H2SO4 soln. of (NH2OH)2·H2SO4. This method prevents problems in the treatment of wastewaters in the clarification area, and eliminates use of addnl. chems. A 50-mm diam., 1.5-m high glass tube was packed with a crosslinked, sulfonic acid group-contg. polystyrene **ion-exchange** resin that was then activated with 5-mol% H2SO4. Next, 116 L wastewater from NH2OH manuf., contg. NH3OH 1.45, H2SO4 0.2, and (NH4)2SO4 0.5 g/L was passed over the resin at 2500 mL/h. Thereafter, the resin was treated with 6500 mL 10-mol% H2SO4, and washed with .apprx.3000 mL water, to obtain 9550 mL (NH2OH)2·H2SO4 soln. contg. NH2OH 19.45, H2SO4 48.06, and (NH4)2SO4 6.7 g/L. The soln. was used in the synthesis of NH2OH.

IT ~~10102-43-9P~~, Nitrogen monoxide, reactions
(hydrogenation of, catalytic, for hydroxylammonium sulfate prepn.,
wastewater treatment in)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

N==O

IC ICM C02F001-42

ICS C01B021-14

CC 49-3 (Industrial Inorganic Chemicals)

Section cross-reference(s): 61

ST hydroxylamine recovery wastewater **ion exchange**;

hydroxylammonium sulfate prepn **ion exchange**;

sulfonated crosslinked polystyrene **ion exchange**

IT **Ion exchangers**

(acidic, in hydroxylamine and hydroxylammonium salt recovery from
wastewater)

IT Wastewater treatment

(**ion exchange**, hydroxylamine and hydroxylamine
salt recovery in)

IT ~~10102-43-9P~~, Nitrogen monoxide, reactions

(hydrogenation of, catalytic, for hydroxylammonium sulfate prepn.,
wastewater treatment in)

IT 9003-53-6D, Polystyrene, crosslinked, sulfonated

(**ion-exchange** resin, in hydroxylamine and
hydroxylammonium salt recovery from wastewaters)

OSC.G 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

L78 ANSWER 10 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 105:122681 HCA Full-text

OREF 105:19711a,19714a

TI Development and active demonstration of acid digestion of plutonium-bearing combustible solid waste

AU Wieczorek, H.; Oser, B.

CS INE, Fed. Rep. Ger.

SO KFK-Nachrichten (~~1986~~), 18(2), 77-82

CODEN: KFKNAW; ISSN: 0340-756X

DT Journal

LA German

AB With the wet-ashing (acid digestion) of .apprx.800 kg of waste and the recovery of 6.3 kg of Pu in a semi-industrial facility, the suitability of the process and the plant components for the treatment of combustible high Pu-contg. wastes is shown. With a suitable reactor constructed for this process, high exchange-rates for the waste and Pu were accomplished. The official requirements were met by the attained decontamination factors for purified off gas of 1010 and for the liq. secondary waste of >106. For 1 kg of wet-ashed waste, 2.3 kg of secondary waste were obtained.

IT ~~9002-88-4~~

(acid digestion of combustible solid waste contg.)

RN 9002-88-4 HCA

CN Ethene, homopolymer (CA INDEX NAME)

CM 1

CRN 74-85-1

CMF C2 H4



IT ~~10102-43-9P~~, preparation

(formation of, in wet-ashing of combustible plutonium-solid wastes)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



CC 71-11 (Nuclear Technology)

IT 7782-50-5D, compds. 9002-86-2 ~~9002-88-4~~ 7440-07-5, uses and miscellaneous

(acid digestion of combustible solid waste contg.)

IT 7446-09-5P, preparation 7647-01-0P, preparation ~~10102-43-9P~~, preparation

(formation of, in wet-ashing of combustible plutonium-solid wastes)
OSC.G 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L78 ANSWER 11 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 105:98084 HCA Full-text

OREF 105:15875a,15878a

TI Derivatization reactions on oxidized polyolefins

AU Carlsson, D. J.; Brousseau, R.; Zhang, Can; Wiles, D. M.

CS Div. Chem., Natl. Res. Counc. Canada, Ottawa, ON, K1A 0R9, Can.

SO Polymer Preprints (American Chemical Society, Division of Polymer Chemistry) (1986), 27(2), 97-8

CODEN: ACPPAY; ISSN: 0032-3934

DT Journal

LA English

AB Polypropylene and LDPE could be smoothly oxidized by α -irradn. The hydroperoxide groups resulting from this oxidn. were extremely reactive to several gaseous reagents at room temp. and could be converted to fluorides, hydrosulfates, alkyl peroxides, chloroformates, and nitrates.

IT 9002-88-4DP, oxidized, derivs. 10102-43-9DP, reaction products with oxidized polypropylene and LDPE (prepn. and characterization of)


RN 9002-88-4 HCA

CN Ethene, homopolymer (CA INDEX NAME)

CM 1

CRN 74-85-1

CMF C2 H4



RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



CC 35-8 (Chemistry of Synthetic High Polymers)

IT 75-44-5DP, reaction products with oxidized polypropylene and LDPE

334-88-3DP, reaction products with oxidized polypropylene and LDPE

7446-09-5DP, reaction products with oxidized polypropylene and LDPE

7783-60-0DP, reaction products with oxidized polypropylene and LDPE

9002-88-4DP, oxidized, derivs. 9003-07-0DP, oxidized,

derivs. 10102-43-9DP, reaction products with oxidized

polypropylene and LDPE

(prepn. and characterization of)

L78 ANSWER 12 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 98:42709 HCA Full-text

OREF 98:6475a,6478a

TI Formation of radiolytically induced gases from solid products of low-level and intermediate-level radioactive wastes

AU Schorr, W.; Duschner, H.; Starke, K.

CS Kernchem. Fachber. Phys. Chem., Philips-Univ. Marburg, Marburg/Lahn, D-3550, Fed. Rep. Ger.

SO Nukleare Entsorgung (1981), 1, 263-75
CODEN: NUKEDA; ISSN: 0723-0893

DT Journal

LA German

AB In the org. material studied, the principal components of the radiolytic gases produced are formed by radiolytically induced chain reactions. Thus H is formed in bitumen and polyethylene. The rate of formation is slow and practically independent of dose rate but linearly dependent on total dose. This relation holds over the dose range expected from fission products whose sp. radioactivity 2 yr after removal from the reactor is 0.1-1 Ci/L. Only small amts. of addnl. gas are formed by admixt. of simulated wastes. The prodn. rate of H in pure matrix remains unaltered. Because O is adsorbed onto the surface of the org. material, the atm. surrounding the waste containers strongly depends on the design of the storage chamber. In contrast to the org. matrixes gas formation in concrete is influenced by such admixts. The detn. of the qual. and quant. compn. of multicomponent gas mixts. was carried out using mass spectroscopy. The complex mass spectra obtained are subjected to math. anal. followed by statistical methods of error redn.

IT 10102-43-98, preparation

(formation of, from irradiated solid products of low-level and intermediate-level radioactive waste)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)

$\text{N}=\text{O}$

IT 9002-88-4

(hydrogen formation by radiolysis of radioactive waste contg.)

RN 9002-88-4 HCA

CN Ethene, homopolymer (CA INDEX NAME)

CM 1

CRN 74-85-1

CMF C2 H4

$\text{H}_2\text{C}=\text{CH}_2$

CC 71-11 (Nuclear Technology)
Section cross-reference(s): 58
IT Cement
 Ion exchangers
 Surfactants
 (formation of gases from radioactive wastes contg.)
IT 74-89-5P, preparation 75-50-3P, preparation 630-08-0P, preparation
7782-44-7P, preparation 10102-43-9P, preparation
10102-44-0P, preparation
 (formation of, from irradiated solid products of low-level and
 intermediate-level radioactive waste)
IT 9002-88-4
 (hydrogen formation by radiolysis of radioactive waste contg.)
OSC.G 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L78 ANSWER 13 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 87:185330 HCA Full-text

OREF 87:29289a,29292a

TI Measurement of toxic substances in the combustion products of certain
construction plastics

AU Oksanen, Pekka; Kallonen, Raija

CS Finland

SO Palontorjuntatekniikka (1975), (2), 48-50

CODEN: PALODT; ISSN: 0031-0476

DT Journal

LA Finnish

AB The rates of formation of CO, HCN, HCl, NO, and NO2 in the combustion of
various plastics were detd. in a smoke density chamber by subjecting test
samples to heat radiation of 2.5W/cm2. Most plastics were less hazardous
than pine wood during combustion. Low-density polyethylene [9002-88-4] had
the highest rate of formation of CO, whereas formation of HCl was fastest in
PVC [9002-86-2]. The highest toxicity indexes belonged to a phenolic foam
(due to CO), a polyurethane foam (due mainly to HCN and NO2), and PVC (due
mainly to HCl).

IT 9002-88-4
 (combustion products of, toxicity of)

RN 9002-88-4 HCA

CN Ethene, homopolymer (CA INDEX NAME)

CM 1

CRN 74-85-1

CMF C2 H4

$\text{H}_2\text{C}=\text{CH}_2$

IT 10102-43-9P, preparation
 (formation of, in combustion of plastic building materials)

RN 10102-43-9 HCA
CN Nitrogen oxide (NO) (CA INDEX NAME)



CC 36-4 (Plastics Manufacture and Processing)
IT 9002-86-2 ~~9002-88-4~~
(combustion products of, toxicity of)
IT 74-90-8P, preparation 630-08-0P, preparation 7647-01-0P,
preparation ~~10102-43-9P~~, preparation 10102-44-0P,
preparation
(formation of, in combustion of plastic building materials)

L78 ANSWER 14 OF 14 HCA COPYRIGHT 2010 ACS on STN

AN 57:84783 HCA Full-text

OREF 57:16985b-d

TI Use of ~~ion-exchanging~~ resins for purification of
nonvolatile aliphatic acids by paper chromatography

AU Fateeva, M. V.

SO Biokhimiya (Moscow) (~~1962~~), 27, 32-7

CODEN: BIOHAO; ISSN: 0320-9725

DT Journal

LA Unavailable

AB An acid-contg. soln. (100 ml.) was passed through a cationite SDV-3 (50 ml./hr.) until a blue color developed in the bromocresol green test. The eluate was immediately passed through the anionite N-O column and the absence of sugar was checked. To remove all the acids (checked by titration with 0.1N HCl against methyl red) and regenerate the column, 100 ml. of 3% NaOH was passed through it. Then the eluate was immediately passed through another column contg. SDV-3. Free acids were collected and examd. chromatographically. Good results were obtained with mixts. contg. sugars, alcs., amino acids, and inorg. salts.

IT ~~10102-43-9P~~, N-O
(in aliphatic-acid purification)

RN 10102-43-9 HCA

CN Nitrogen oxide (NO) (CA INDEX NAME)



CC 55 (Biochemical Methods)

IT ~~Ion exchange~~

(acid purification by)

IT Acids

(catalysts in polymerization, purification of aliphatic,
~~ion-exchanging~~ resins in)

IT ~~10102-43-9P~~, N-O 12778-16-4P, SDV 3

(in aliphatic-acid purification)